

Anti-Lamin B1 antibody [119D5-F1] - Nuclear Envelope Marker ab8982

KO 評価済

★★★★☆ 6 Abreviews 40 References 画像数 7

製品の概要

| | |
|--------------|---|
| 製品名 | Anti-Lamin B1 antibody [119D5-F1] - Nuclear Envelope Marker |
| 製品の詳細 | Mouse monoclonal [119D5-F1] to Lamin B1 - Nuclear Envelope Marker |
| 由来種 | Mouse |
| 特異性 | Reacts with an epitope located C-terminal of residue 231 in lamin B1. Reacts against lamin B1, does not cross react with lamin B2. Lamins do not appear to be universally distributed among different cell and tissue types. ab8982 has been shown to react with HeLa and 3T3 cells in immunocytochemistry. Other cell/tissue types have not been tested. |
| アプリケーション | 適用あり: Flow Cyt, ICC/IF, IHC-Fr |
| 種交差性 | 交差種: Mouse, Human, Pig |
| 免疫原 | Full length native protein (purified) corresponding to Rat Lamin B1 (C terminal). Immunogen is purified rat liver lamins. Database link: P20700 |
| エピトープ | C-terminal to residue 231. |
| ポジティブ・コントロール | ICC/IF: HAP1, HeLa and 3T3 cells. WB: Human mammary cell lysate. IHC-Fr: Human kidney and colon tissue; MCF7 cell culture; Swine liver tissue. Flow Cyt: HeLa cells |
| 特記事項 | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

製品の特性

| | |
|-------|---|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. |

| | |
|--------|--|
| バッファー | Preservative: 0.09% Sodium azide Constituent: 99% PBS |
| 精製度 | Protein G purified |
| ポリ/モノ | モノクローナル |
| クローン名 | 119D5-F1 |
| ミエローマ | P3x63-Ag8.653 |
| アイソタイプ | IgG1 |
| 軽鎖の種類 | kappa |

アプリケーション


The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab8982の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|--|
| Flow Cyt | | 1/100 - 1/200. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody. |
| ICC/IF | | Use a concentration of 0.5 µg/ml. |
| IHC-Fr | | Use at an assay dependent concentration. |

ターゲット情報

| | |
|-------|--|
| 機能 | Lamins are components of the nuclear lamina, a fibrous layer on the nucleoplasmic side of the inner nuclear membrane, which is thought to provide a framework for the nuclear envelope and may also interact with chromatin. |
| 関連疾患 | Defects in LMNB1 are the cause of leukodystrophy demyelinating autosomal dominant adult-onset (ADLD) [MIM:169500]. ADLD is a slowly progressive and fatal demyelinating leukodystrophy, presenting in the fourth or fifth decade of life. Clinically characterized by early autonomic abnormalities, pyramidal and cerebellar dysfunction, and symmetric demyelination of the CNS. It differs from multiple sclerosis and other demyelinating disorders in that neuropathology shows preservation of oligodendroglia in the presence of subtotal demyelination and lack of astrogliosis. |
| 配列類似性 | Belongs to the intermediate filament family. |
| 翻訳後修飾 | B-type lamins undergo a series of modifications, such as farnesylation and phosphorylation. Increased phosphorylation of the lamins occurs before envelope disintegration and probably plays a role in regulating lamin associations. |
| 細胞内局在 | Nucleus inner membrane. |


画像



Immunocytochemistry/ Immunofluorescence - Anti-Lamin B1 antibody [119D5-F1] - Nuclear Envelope Marker (ab8982)


ab8982 staining Lamin B1 in wild-type HAP1 cells (top panel) and LMNB1 knockout HAP1 cells (bottom panel). The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab8982 at 0.5µg/ml and **ab202272** at 1/250 dilution (shown in pseudocolour red) overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to Mouse IgG (Alexa Fluor® 488) (**ab150117**) at 2 µg/ml (shown in green). Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunohistochemistry (Frozen sections) - Anti-Lamin B1 antibody [119D5-F1] - Nuclear Envelope Marker (ab8982)

Immunohistochemistry on frozen sections of human kidney showing nuclear lamina staining in epithelial and connective tissue cells.



Immunohistochemistry (Frozen sections) - Anti-Lamin B1 antibody [119D5-F1] - Nuclear Envelope Marker (ab8982)

Immunohistochemistry of MCF-7 cell culture showing nuclear lamina staining.



Immunocytochemistry/ Immunofluorescence - Anti-Lamin B1 antibody [119D5-F1] - Nuclear Envelope Marker (ab8982)

This image is courtesy of Marilena Ciciarello & Patrizia Lavia, University La Sapienza


Methanol fixed HeLa and 3T3 cells ([ab7179](#)) were stained with ab8982 (1/50 and 1/20 respectively). The cells were fixed in 100% methanol for 6 minutes at -20°C. ab8982 clearly stains the nuclear envelope with very little background staining.

A: HeLa cells + ab8982 (1/50) (green)

B: HeLa cells counterstained with DAPI (blue)

C: 3T3 cells + ab8982 (1/20) (green)


D: 3T3 cells counterstained with DAPI (blue)



Flow Cytometry - Anti-Lamin B1 antibody [119D5-F1] - Nuclear Envelope Marker (ab8982)


Overlay histogram showing HeLa cells stained with ab8982 (red line). The cells were fixed with 100% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum ([ab7481](#)) / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab8982, 1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was Mouse IgG1 [ICIGG1] ([ab91353](#), 2µg/1x10⁶ cells) used under the same conditions.

Acquisition of >5,000 events was performed.



Immunohistochemistry on frozen sections of swine liver showing nuclear lamina staining in hepatocytes.

Immunohistochemistry (Frozen sections) - Anti-Lamin B1 antibody [119D5-F1] - Nuclear Envelope Marker (ab8982)



Immunohistochemistry on frozen sections of human colon showing nuclear lamina staining in epithelial and connective tissue cells.

Immunohistochemistry (Frozen sections) - Anti-Lamin B1 antibody [119D5-F1] - Nuclear Envelope Marker (ab8982)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors