

Anti-Lipocalin-2 / NGAL antibody ab89819

画像数 3

製品の概要

製品名	Anti-Lipocalin-2 / NGAL antibody
製品の詳細	Mouse polyclonal to Lipocalin-2 / NGAL
由来種	Mouse
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human
免疫原	Recombinant full length protein within Human Lipocalin-2/ NGAL aa 1-200. The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please <u>contact</u> our Scientific Support team to discuss your requirements. Database link: <u>NP_005555.1</u>
ポジティブ・コントロール	WB: spleen tissue lysate. IHC: pancreas tissue.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.40 Constituent: 100% PBS
精製度	Protein A purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab89819の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		
IHC-P		

追加情報

IHC-P: Use at a concentration of 3 µg/ml. Antigen retrieval is not essential but may optimise staining. Microwave method recommended.

WB: Use at a concentration of 1 µg/ml. Predicted molecular weight: 23 kDa.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

機能

Iron-trafficking protein involved in multiple processes such as apoptosis, innate immunity and renal development. Binds iron through association with 2,5-dihydroxybenzoic acid (2,5-DHBA), a siderophore that shares structural similarities with bacterial enterobactin, and delivers or removes iron from the cell, depending on the context. Iron-bound form (holo-24p3) is internalized following binding to the SLC22A17 (24p3R) receptor, leading to release of iron and subsequent increase of intracellular iron concentration. In contrast, association of the iron-free form (apo-24p3) with the SLC22A17 (24p3R) receptor is followed by association with an intracellular siderophore, iron chelation and iron transfer to the extracellular medium, thereby reducing intracellular iron concentration. Involved in apoptosis due to interleukin-3 (IL3) deprivation: iron-loaded form increases intracellular iron concentration without promoting apoptosis, while iron-free form decreases intracellular iron levels, inducing expression of the proapoptotic protein BCL2L11/BIM, resulting in apoptosis. Involved in innate immunity, possibly by sequestering iron, leading to limit bacterial growth.

組織特異性

Expressed in bone marrow and in tissues that are prone to exposure to microorganism. High expression is found in bone marrow as well as in uterus, prostate, salivary gland, stomach, appendix, colon, trachea and lung. Not found in the small intestine or peripheral blood leukocytes.


配列類似性

Belongs to the calycin superfamily. Lipocalin family.

細胞内局在

Secreted. Upon binding to the SLC22A17 (24p3R) receptor, it is internalized.

画像




Western blot - Anti-Lipocalin-2 / NGAL antibody
(ab89819)

Anti-Lipocalin-2 / NGAL antibody (ab89819) at 1 µg/ml + Human spleen tissue lysate at 50 µg

Predicted band size: 23 kDa

Observed band size: 23 kDa



Western blot - Anti-Lipocalin-2 / NGAL antibody
(ab89819)

All lanes : Anti-Lipocalin-2 / NGAL antibody (ab89819) at 1 µg/ml

Lane 1 : Lipocalin 2 transfected 293T cell lysate


Lane 2 : Non-transfected lysate

Lysates/proteins at 25 µg per lane.

Predicted band size: 23 kDa

Observed band size: 23 kDa

Additional bands at: ~60 kDa. We are unsure as to the identity of these extra bands.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lipocalin-2 / NGAL antibody (ab89819)

ab89819 at 3µg/ml staining Lipocalin-2/ NGAL in formalin-fixed, paraffin-embedded Human pancreas tissue by immunohistochemistry .

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors