

# Anti-Listeria monocytogenes antibody ab35132

★★★★★ [1 Abreviews](#) [17 References](#)

### 製品の概要

---

製品名	Anti-Listeria monocytogenes antibody
製品の詳細	Rabbit polyclonal to Listeria monocytogenes
由来種	Rabbit
特異性	This antibody is specific for Listeria monocytogenes whole cells. This antiserum has not been absorbed and may react with related microorganisms.
アプリケーション	<b>適用あり:</b> WB, ELISA, IHC-P, ICC/IF
種交差性	<b>交差種:</b> Listeria monocytogenes
免疫原	Tissue, cells or virus corresponding to Listeria monocytogenes. Listeria monocytogenes. whole cells. ATCC strain 43251.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

---

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.20 Preservative: 0.1% Sodium azide Constituent: PBS
特記事項(精製)	>95% pure.
ポリモノ	ポリクローナル
アイソタイプ	IgG

### アプリケーション

---

## The Abpromise guarantee

**Abpromise保証は、次のテスト済みアプリケーションにおけるab35132の使用に適用されます**

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		
ELISA		
IHC-P		
ICC/IF		

### 追加情報

ELISA: 1/500 - 1/2000.

ICC/IF: Use at an assay dependent dilution.

IHC-P: Use at an assay dependent dilution. (Pre-treatment with 0.1% pronase was used)

WB: Use at an assay dependent dilution.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

### ターゲット情報

#### 関連性

The genus *Listeria* comprises six species: *L. monocytogenes*, *L. innocua*, *L. welshimeri*, *L. seeligeri*, *L. ivanovii* and *L. grayi*. *Listeria monocytogenes*, the most commonly isolated pathogenic member, is associated with a wide spectrum of human and animal diseases. In the smear from the original tissue, *L. monocytogenes* may appear as gram-positive coccobacilli that may be confused with *Streptococcus agalactiae* (group B), enterococci, or *Corynebacterium* spp. *Listeria* is differentiated from streptococci by a positive catalase test. *L. monocytogenes* is the only species of the genus *Listeria* that has been clearly documented as a pathogen for humans. The forms of disease caused by this organism are myriad and age-related. The most common clinical manifestations are meningitis and septicemia. *Listeria monocytogenes*, a food-borne intracellular animal and human pathogen, interacts with infected host cells both prior to entry and during the intracellular phase of infection.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors