


Anti-MAL2 antibody ab75347

8 References

製品の概要

製品名	Anti-MAL2 antibody
製品の詳細	Rabbit polyclonal to MAL2
由来種	Rabbit
アプリケーション	適用あり: ELISA, ICC, WB
種交差性	交差種: Human 交差が予測される動物種: Mouse, Rat 
免疫原	Synthetic peptide derived from the N-terminal domain of human MAL2
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	Constituent: Whole serum
精製度	Whole antiserum
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab75347の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ELISA		Use at an assay dependent concentration.
ICC		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration.

ターゲット情報

機能	Member of the machinery of polarized transport. Required for the indirect transcytotic route at the step of the egress of the transcytosing cargo from perinuclear endosomes in order for it to travel to the apical surface via a raft-dependent pathway.
組織特異性	Predominantly expressed in kidney, lung, and liver. Also found in thyroid gland, stomach and, at lower levels in testis and small intestine.
配列類似性	Belongs to the MAL family. Contains 1 MARVEL domain.
細胞内局在	Cell membrane. Apical cell membrane. Endomembrane system. Cytoplasm > perinuclear region. Associated with lipid rafts. In polarized epithelial cells, restricted to the apical surface. In hepatocytes, as well as in polarized hepatoma HepG2 cells, found in the canalicular membrane, equivalent to the apical surface, beneath the canalicular actin cytoskeleton. In non-polarized HepG2 cells, distributed to the perinuclear region.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors