


Anti-MOX1/MEOX1 antibody ab125712

画像数 3

製品の概要

製品名	Anti-MOX1/MEOX1 antibody
製品の詳細	Rabbit polyclonal to MOX1/MEOX1
由来種	Rabbit
アプリケーション	適用あり: WB
種交差性	交差種: Human 交差が予測される動物種: Chimpanzee, Macaque monkey, Gorilla, Orangutan 
免疫原	Synthetic peptide within Human MOX1/MEOX1 aa 50-150 conjugated to keyhole limpet haemocyanin. The exact sequence is proprietary. Database link: P50221
ポジティブ・コントロール	This antibody gave a positive signal in Human Fetal Heart tissue lysate. WB: Human liver cell lysate
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 2% Bovine Serum Albumin before being incubated with abX overnight at 4°C. Antibody binding was detected using an anti-rabbit

	antibody conjugated to HRP, and visualised using ECL development solution ab133406
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab125712の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 34 kDa (predicted molecular weight: 27 kDa).

ターゲット情報

機能	Mesodermal transcription factor that plays a key role in somitogenesis and is specifically required for sclerotome development. Required for maintenance of the sclerotome polarity and formation of the cranio-cervical joints (PubMed:23290072, PubMed:24073994). Binds specifically to the promoter of target genes and regulates their expression. Activates expression of NKX3-2 in the sclerotome. Activates expression of CDKN1A and CDKN2A in endothelial cells, acting as a regulator of vascular cell proliferation. While it activates CDKN1A in a DNA-dependent manner, it activates CDKN2A in a DNA-independent manner. Required for hematopoietic stem cell (HSCs) induction via its role in somitogenesis: specification of HSCs occurs via the deployment of a specific endothelial precursor population, which arises within a sub-compartment of the somite named endotome.
関連疾患	Klippel-Feil syndrome 2, autosomal recessive
配列類似性	Contains 1 homeobox DNA-binding domain.
細胞内局在	Nucleus. Cytoplasm. Localizes predominantly in the nucleus.

画像



Western blot - Anti-MOX1/MEOX1 antibody
(ab125712)

All lanes : Anti-MOX1/MEOX1 antibody (ab125712) at 1 µg/ml

Lane 1 : Human liver cell lysate

Lane 2 : MCF7 cell lysate


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 27 kDa

Observed band size: 28 kDa

False colour image of Western blot: Anti-MOX1/MEOX1 antibody staining at 1 ug/ml, shown in green; Mouse anti-Alpha Tubulin [DM1A] ([ab7291](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab125712 was shown to bind specifically to MOX1/MEOX1. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in fluorescent western blot (TBS-based) blocking solution before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ([ab216776](#)) at 1/20000 dilution.



Western blot - Anti-MOX1/MEOX1 antibody
(ab125712)

Anti-MOX1/MEOX1 antibody (ab125712) at 1 µg/ml + Heart
(Human) Whole Cell Lysate - fetal normal tissue at 25 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/50000 dilution

Developed using the ECL technique.

Performed under reducing conditions.


Predicted band size: 27 kDa

Observed band size: 34 kDa

Additional bands at: 65 kDa (possible non-specific binding), 75 kDa (possible non-specific binding)

Exposure time: 20 minutes

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 2% Bovine Serum Albumin before being incubated with ab125712 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution **ab133406**



All lanes : Anti-MOX1/MEOX1 antibody (ab125712) at 1 µg/ml

Lane 1 : Heart (Human) Whole Cell Lysate - fetal normal tissue

Lane 2 : Heart (Human) Whole Cell Lysate - fetal normal tissue with Immunising peptide at 1 µg/ml

Lysates/proteins at 25 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/50000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 27 kDa

Observed band size: 34 kDa

Additional bands at: 30 kDa (possible non-specific binding), 65 kDa (possible non-specific binding)

Exposure time: 20 minutes

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 2% Bovine Serum Albumin before being incubated with ab125712 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody

conjugated to HRP, and visualised using ECL development solution

ab133406

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors