


### Anti-MTCO1 antibody [1D6E1A8] ab14705

リコンビナント

★★★★★ **42 Abreviews**   **474 References**   画像数 12

#### 製品の概要

製品名	Anti-MTCO1 antibody [1D6E1A8]
製品の詳細	Mouse monoclonal [1D6E1A8] to MTCO1
由来種	Mouse
アプリケーション	<b>適用あり:</b> ICC, IHC-P, WB, Flow Cyt
種交差性	<b>交差種:</b> Mouse, Rat, Human <b>交差が予測される動物種:</b> Sheep, Goat, Cat, Dog, Pig, Caenorhabditis elegans, Zebrafish, Quail, Rhesus monkey, Chinese hamster, Common marmoset 
免疫原	Full length native protein (purified). This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: HeLa, MCF7, C6, PC-12, Neuro-2a, RAW 264.7 whole cell lysates. Human skeletal, Rat heart and Rat liver tissue lysates. IHC-P: Human kidney, Mouse colon, and Rat colon tissue. ICC: Human cervical adenocarcinoma, Mouse neuroblastoma neuroblast, and rat glial tumor cells. Flow Cyt: HeLa, Neuro-2a, and C6 cells.
特記事項	This product has switched from a hybridoma to recombinant production method on 24th May 2023.

#### **Western blot protocol advice:**

For best results with this antibody in Western blot, do not boil samples before loading onto the gel. Boiling of the sample will cause a loss of signal.

Hydrophobic intrinsic membrane proteins such as the core mtDNA-encoded proteins of the mitochondrial OXPHOS complexes tend to run faster in SDS-PAGE than predicted by their amino acid composition. This is likely due to incomplete unfolding of the protein and a more negative charge:mass ratio.

This antibody clone [1D6E1A8] is manufactured by Abcam. If you require a different buffer formulation or a particular conjugate for your experiments, please contact [orders@abcam.com](mailto:orders@abcam.com).

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

## 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	1D6E1A8
アイソタイプ	IgG2a
軽鎖の種類	kappa

## アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、**次のテスト済みアプリケーションにおけるab14705の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC	★★★★★ (2)	1/100.
IHC-P	★★★★★ (13)	1/5000.
WB	★★★★★ (18)	1/1000.
Flow Cyt	★★★★★ (1)	1/500.

## ターゲット情報

機能	Cytochrome c oxidase is the component of the respiratory chain that catalyzes the reduction of oxygen to water. Subunits 1-3 form the functional core of the enzyme complex. CO I is the catalytic subunit of the enzyme. Electrons originating in cytochrome c are transferred via the copper A center of subunit 2 and heme A of subunit 1 to the bimetallic center formed by heme A3 and copper B.
パスウェイ	Energy metabolism; oxidative phosphorylation.
関連疾患	Defects in MT-CO1 are a cause of Leber hereditary optic neuropathy (LHON) [MIM:535000]. LHON is a maternally inherited disease resulting in acute or subacute loss of central vision, due to optic nerve dysfunction. Cardiac conduction defects and neurological defects have also been described in some patients. LHON results from primary mitochondrial DNA mutations affecting the respiratory chain complexes. Defects in MT-CO1 are a cause of anemia sideroblastic acquired idiopathic (AISA) [MIM:516030]; a disease characterized by inadequate formation of heme and excessive accumulation of iron in mitochondria. Defects in MT-CO1 are a cause of mitochondrial complex IV deficiency (MT-C4D) [MIM:220110]; also known as cytochrome c oxidase deficiency. A disorder of the mitochondrial respiratory chain

with heterogeneous clinical manifestations, ranging from isolated myopathy to severe multisystem disease affecting several tissues and organs. Features include hypertrophic cardiomyopathy, hepatomegaly and liver dysfunction, hypotonia, muscle weakness, exercise intolerance, developmental delay, delayed motor development and mental retardation. A subset of patients manifest Leigh syndrome.

Defects in MT-CO1 are associated with recurrent myoglobinuria mitochondrial (RM-MT) [MIM:550500]. Recurrent myoglobinuria is characterized by recurrent attacks of rhabdomyolysis (necrosis or disintegration of skeletal muscle) associated with muscle pain and weakness, and followed by excretion of myoglobin in the urine.

Defects in MT-CO1 are a cause of deafness sensorineural mitochondrial (DFNM) [MIM:500008]. DFNM is a form of non-syndromic deafness with maternal inheritance. Affected individuals manifest progressive, postlingual, sensorineural hearing loss involving high frequencies. Defects in MT-CO1 are a cause of colorectal cancer (CRC) [MIM:114500].


## 配列類似性

Belongs to the heme-copper respiratory oxidase family.

## 細胞内局在

Mitochondrion inner membrane.

## 画像



**All lanes** : Anti-MTCO1 antibody [1D6E1A8] (ab14705) at 1/1000 dilution

**Lane 1** : HeLa (human cervical adenocarcinoma epithelial cell) non-mitochondrial fraction

**Lane 2** : HeLa mitochondrial fraction

**Lane 3** : MCF7 (human breast adenocarcinoma epithelial cell) non-mitochondrial fraction

**Lane 4** : MCF7 mitochondrial fraction

Lysates/proteins at 20 µg per lane.


## Secondary

**All lanes** : Peroxidase-Conjugated Goat anti-Mouse IgG (H+L) at 1/10000 dilution

**Observed band size:** 35 kDa

**Exposure time:** 5 seconds

Blocking and diluting buffer and concentration: 5% NFDM/TBST.



**All lanes :** Anti-MTCO1 antibody [1D6E1A8] (ab14705) at 1/1000 dilution

**Lane 1 :** C6 (rat glial tumor glial cell) whole cell lysate

**Lane 2 :** PC-12 (rat adrenal gland pheochromocytoma cell) whole cell lysate

**Lane 3 :** Neuro-2a (mouse neuroblastoma neuroblast) whole cell lysate

**Lane 4 :** RAW 264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysate

**Lane 5 :** Human skeletal muscle tissue lysate

**Lane 6 :** Rat heart tissue lysate

**Lane 7 :** Rat liver tissue lysate


Lysates/proteins at 20 µg per lane.

## Secondary


**All lanes :** Peroxidase-Conjugated Goat anti-Mouse IgG (H+L) at 1/10000 dilution

**Observed band size:** 35 kDa

Blocking and diluting buffer and concentration: 5% NFDM/TBST.




Immunohistochemical analysis of paraffin-embedded Human kidney tissue labelling MTCO1 with ab14705 at 1/5000 dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on Human kidney. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, epitope retrieval Solution2) for 20 mins. Counterstained with hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MTCO1 antibody [1D6E1A8] (ab14705)


Immunohistochemical analysis of paraffin-embedded Mouse colon tissue labelling MTCO1 with ab14705 at 1/5000 dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on Mouse colon. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, epitope retrieval Solution2) for 20 mins. Counterstained with hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MTCO1 antibody [1D6E1A8] (ab14705)


Immunohistochemical analysis of paraffin-embedded Rat colon tissue labelling MTCO1 with ab14705 at 1/5000 dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on Rat colon. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, epitope retrieval Solution2) for 20 mins. Counterstained with hematoxylin.






Immunocytochemistry/ Immunofluorescence - Anti-MTCO1 antibody [1D6E1A8] (ab14705)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Human cervical adenocarcinoma epithelial cells labeling MTCO1 with ab14705 at 1/100 dilution, followed by **ab150113** Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 dilution. Confocal image showing cytoplasmic and membranous staining in subsets of Human cervical adenocarcinoma epithelial cells. **ab186735** Anti-TOMM20 antibody [EPR15581-54] - Mitochondrial Marker was used to counterstain tubulin at 1/400 dilution. The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is **ab150080** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 594) preadsorbed at 1/1000 dilution.




Immunocytochemistry/ Immunofluorescence - Anti-MTCO1 antibody [1D6E1A8] (ab14705)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Mouse neuroblastoma neuroblast cells labeling MTCO1 with ab14705 at 1/100 dilution, followed by **ab150113** Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 dilution. Confocal image showing cytoplasmic and membranous staining in subsets of Mouse neuroblastoma neuroblast cells. **ab186735** Anti-TOMM20 antibody [EPR15581-54] - Mitochondrial Marker was used to counterstain tubulin at 1/400 dilution. The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is **ab150080** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 594) preadsorbed at 1/1000 dilution.




Immunocytochemistry/ Immunofluorescence - Anti-MTCO1 antibody [1D6E1A8] (ab14705)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Rat glial tumor cells labeling MTCO1 with ab14705 at 1/100 dilution, followed by **ab150113** Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 dilution. Confocal image showing cytoplasmic and membranous staining in subsets of Rat glial tumor cells. **ab186735** Anti-TOMM20 antibody [EPR15581-54] - Mitochondrial Marker was used to counterstain tubulin at 1/400 dilution. The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is **ab150080** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 594) preadsorbed at 1/1000 dilution.




Flow cytometric analysis of Human HeLa cells labelling MTCO1 with ab14705 at 1/500 dilution (Right) compared with an isotype control (Left). Goat Anti-Rabbit IgG (Alexa Fluor® 488, **ab150081**) at was used as the secondary antibody. Gated on viable cells.

Flow Cytometry - Anti-MTCO1 antibody [1D6E1A8]  
(ab14705)




Flow cytometric analysis of Neuro-2a cells labelling MTCO1 with ab14705 at 1/500 dilution (Right) compared with an isotype control (Left). Goat Anti-Rabbit IgG (Alexa Fluor® 488, **ab150081**) at was used as the secondary antibody. Gated on viable cells.

Flow Cytometry - Anti-MTCO1 antibody [1D6E1A8]  
(ab14705)



Flow cytometric analysis of C6 cells labelling MTCO1 with ab14705 at 1/500 dilution (Right) compared with an isotype control (Left). Goat Anti-Rabbit IgG (Alexa Fluor® 488, **ab150081**) at was used as the secondary antibody. Gated on viable cells.

Flow Cytometry - Anti-MTCO1 antibody [1D6E1A8]  
(ab14705)



**This image was generated using a previous batch manufactured using hybridoma production method.**

250 µm human cerebellar sections from control individuals and a patient with mitochondrial disease underwent passive clearing at 37 °C for 2 or 4 weeks.

The quality of immunofluorescent staining is determined by duration of passive clearing; 2 weeks of passive clearing produced minimal labelling of the white matter in the granule cell layer (NF-H; green; 488 nm and MBP; red, 546 nm) with an absence of labelling of mitochondria (MTCO1 (COXI) (ab14705, 1/100); purple; 647 nm; Extending passive clearing to 4 weeks improved the quality of stain with identifiable Purkinje cells and their axons (NF-H, green; 488 nm) and their myelin sheaths (MBP; red, 546 nm) and mitochondria (MTCO1 (COXI) (ab14705, 1/100); purple; 647 nm.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors