




### Anti-MURF2 antibody ab4387

★★★★★ [2 Abreviews](#) [6 References](#) [画像数 1](#)

#### 製品の概要

製品名	Anti-MURF2 antibody
製品の詳細	Goat polyclonal to MURF2
由来種	Goat
アプリケーション	<b>適用あり:</b> WB, IHC-P
種交差性	<b>交差種:</b> Human <b>交差が予測される動物種:</b> Rat, Dog, Pig 
免疫原	Synthetic peptide: HIFSFSLNSLNE , corresponding to C terminal amino acids 536-546 of Human MURF2. <div>  <a href="#">Run BLAST with</a>  <a href="#">Run BLAST with</a> </div>
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: Tris buffered saline, 0.5% BSA
精製度	Immunogen affinity purified
特記事項(精製)	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
ポリ/モノ	ポリクローナル

## アプリケーション


**The Abpromise guarantee** Abpromise保証は、 次のテスト済みアプリケーションにおけるab4387の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Predicted molecular weight: 60 kDa.
IHC-P		Use a concentration of 10 µg/ml.

## ターゲット情報

機能	May regulate gene expression and protein turnover in muscle cells.
組織特異性	Highly expressed in muscle. Low-level expression in liver.
配列類似性	Contains 1 B box-type zinc finger. Contains 1 COS domain. Contains 1 RING-type zinc finger.
細胞内局在	Cytoplasm. Nucleus. Nuclear under atrophic conditions and upon mechanical signals. Localizes to the sarcomeric M-band in cardiomyocytes. Colocalizes in part with microtubules.

## 画像



Anti-MURF2 antibody (ab4387) at 0.1 µg/ml + Human heart lysate  
(in RIPA buffer) at 35 µg

**Predicted band size:** 60 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors