



Anti-Myc tag antibody ab9106

★★★★★ [39 Abreviews](#) [472 References](#) [画像数 7](#)

製品の概要

製品名	Anti-Myc tag antibody
製品の詳細	Rabbit polyclonal to Myc tag
由来種	Rabbit
特異性	ELISA: Antibody specificity was verified by ELISA against the peptide (EQKLISEEDL). A 1:60,000 dilution of the antibody gave an O.D.=1.0 in a 15 minute reaction using HRP-conjugated Goat Anti Rabbit IgG at 1:20,000 and TMB as the substrate. Appropriate specificity controls were run.
アプリケーション	適用あり: IHC-Fr, IP, WB, IHC-P, ICC/IF, Electron Microscopy
種交差性	交差種: Species independent
免疫原	Synthetic peptide corresponding to Myc tag conjugated to keyhole limpet haemocyanin. (Peptide available as ab13837)
ポジティブ・コントロール	<div>  Run BLAST with  Run BLAST with </div> <p>ICC/IF: CHO cells transfected with 12-tags constructs. IHC-P: Mouse brain tissue stained for Myc tag. EM: Mouse embryonic fibroblasts expressing myc-tagged protein. IP: Huh7 expressing myc-tagged CDC42 protein and transfected 293FT cells. WB: Y. enterocolitica ?HOPEMT carrying plasmids encoding Myc-tagged CT082 or CT695, and HA-tagged Slc1.</p>
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.1% Sodium azide
精製度	Affinity purified

特記事項 (精製)	Antibodies were immunoaffinity purified using the peptide immobilized on a solid phase. Antibody concentration was determined by extinction coefficient : O.D. 1.4 at 280nm equals 1.0 mg of IgG.
ポリモノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab9106の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-Fr	★★★★★ (1)	1/500.
IP	★★★★★ (9)	Use at an assay dependent concentration.
WB	★★★★★ (14)	Use at an assay dependent concentration.
IHC-P	★★★★★ (3)	Use at an assay dependent concentration.
ICC/IF	★★★★★ (10)	1/1000.
Electron Microscopy		Use at an assay dependent concentration. See Abreview for further details (submitted by Eeva-Liisa Eskelinen).

ターゲット情報


関連性	Epitope tags are short peptide sequences that are easily recognized by tag-specific antibodies. Due to their small size, epitope tags do not affect the tagged protein's biochemical properties. Most often sequences encoding the epitope tag are included with target DNA at the time of cloning to produce fusion proteins containing the epitope tag sequence. This allows anti-epitope tag antibodies to serve as universal detection reagents for any tag containing protein produced by recombinant means. This means that anti-epitope tag antibodies are a useful alternative to generating specific antibodies to identify, immunoprecipitate or immunoaffinity purify a recombinant protein. The anti-epitope tag antibody is usually functional in a variety of antibody-dependent experimental procedures. Expression vectors producing epitope tag fusion proteins are available for a variety of host expression systems including bacteria, yeast, insect and mammalian cells.
細胞内局在	Nuclear

画像



Immunocytochemistry/ Immunofluorescence - Anti-Myc tag antibody (ab9106)

Immunofluorescent analysis of 4% paraformaldehyde (PFA)-fixed, permeabilized with 0.1% Triton X-100 in CHO cells transfected with 12-tags constructs (named as CHO-12 tags, Top) and parental CHO cells (Bottom) labelling MYC with ab9106 at 1/1000 dilution (1 µg/mL), followed by Donkey Anti-rabbit IgG (H&L), FITC conjugated antibody at 1/1000 dilution (1 µg/mL) (Green). Nucleus was counterstained with DAPI (Blue).



Western blot - Anti-Myc tag antibody (ab9106)

Pais et al PLoS One. 2013;8(2):e56292. doi: 10.1371/journal.pone.0056292. Epub 2013 Feb 19. Fig 6. Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>

Y. enterocolitica ΔHOPEMT carrying plasmids encoding Myc-tagged CT082 or CT695, and HA-tagged Slc1 (as indicated) were grown in non-secreting conditions. Chloramphenicol was added (time=?0 min) to stop bacterial protein synthesis and samples were taken at the indicated time points. Samples were analyzed by immunoblotting using ab9106, anti-HA, and anti-TEM antibodies.




Immunoprecipitation - Anti-Myc tag antibody
(ab9106)

This image is courtesy of an anonymous abreview.

ab9106 at 4 µg/mg lysate used in Huh7 (Human cell line) whole cell lysate (1×10^6 cells).

Cells expressed myc-tagged CDC42 protein. Immunoprecipitation step performed using Protein A matrix. Incubation time 4 hours at 4°C. Western Blot antibody used at 1/300 dilution.



Immunoprecipitation - Anti-Myc tag antibody
(ab9106)

ab9106 used at 1/250 in the immunoprecipitation of transfected human 293FT cells (whole cell lysate).

In this experiment cells were co-transfected with myc tagged cdc25 along with an HA tagged viral protein which binds to cdc25. The cell lysate was immunoprecipitated using ab9106, run on a 10% gel. The blot was probed for HA and HA tagged protein (which co-IPed with Myc tagged protein) and this is seen at ~15KDa.


Lane 1: Input non-transfected cells

Lane 2: Immunoprecipitated non-transfected cells

Lane 3: Input co-transfected cells

Lane 4: Immunoprecipitated co-transfected cells

This image is courtesy of an Abreview submitted on **23 March 2006**.




Immunohistochemistry (Frozen sections) - Anti-Myc tag antibody (ab9106)

This image is courtesy of an anonymous abreview.

Paraformaldehyde-fixed frozen mouse brain tissue stained for Myc tag using ab9106 at a 1/500 dilution (2hrs at 25°C), followed by a goat anti-rabbit Alexa-Fluor® 568 secondary used at a 1/500 dilution.

For further details please see [abreview](#).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Myc tag antibody (ab9106)

This image is courtesy of Karen Lee by Abreview.

Paraformaldehyde-fixed, 0.3% Triton X-100 permeabilized mouse brain tissue stained for Myc tag using ab9106 at a 1/500 dilution (15 hrs, 4°C) followed by and Alexa-Fluor® 555 donkey anti-rabbit secondary used at a 1/500 dilution.

For further details please see [abreview](#).



Electron Microscopy - Anti-Myc tag antibody
(ab9106)

This image is courtesy of an Abreview submitted by Dr
Eeva-Liisa Eskelinen

ab9106 at 1/2000 staining mouse embryonic fibroblasts expressing myc-tagged protein by ICC.

The cells were paraformaldehyde fixed and blocked with 3% BSA prior to incubation with the antibody for 1 hour. A goat anti-rabbit IgG antibody (10 nm gold conjugated) was used as the secondary.

The image shown is immuno electron microscopical staining with thawed cryosections.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors