


Anti-Myeloperoxidase antibody ab45977

★★★★★ 17 Abreviews 74 References 画像数 5

製品の概要

製品名	Anti-Myeloperoxidase antibody
製品の詳細	Rabbit polyclonal to Myeloperoxidase
由来種	Rabbit
特異性	ab45977 has been batch tested using human lysates/tissues only. Some customers have successfully used ab45977 with rat. Please contact Abcam Scientific Support for more information.
アプリケーション	適用あり: ICC/IF, WB, IHC-P
種交差性	交差種: Human 交差が予測される動物種: Rat, Common marmoset 
免疫原	Synthetic peptide corresponding to Human Myeloperoxidase aa 150-250 conjugated to keyhole limpet haemocyanin. (Peptide available as ab45976)
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS
	Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab45977の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF	★★★★★ (1)	Use a concentration of 5 - 10 µg/ml.
WB	★★★★★ (2)	Use a concentration of 1 µg/ml. Detects a band of approximately 84 kDa (predicted molecular weight: 84 kDa).
IHC-P	★★★★★ (11)	Use a concentration of 1 µg/ml.

ターゲット情報


機能	Part of the host defense system of polymorphonuclear leukocytes. It is responsible for microbicidal activity against a wide range of organisms. In the stimulated PMN, MPO catalyzes the production of hypohalous acids, primarily hypochlorous acid in physiologic situations, and other toxic intermediates that greatly enhance PMN microbicidal activity.
関連疾患	Defects in MPO are the cause of myeloperoxidase deficiency (MPD) [MIM:254600]. MPD is an autosomal recessive defect that results in disseminated candidiasis.
配列類似性	Belongs to the peroxidase family. XPO subfamily.
細胞内局在	Lysosome.

画像



Immunocytochemistry/ Immunofluorescence - Anti-Myeloperoxidase antibody (ab45977)


ab45977 stained in HL60 cells. Cells were fixed with 100% methanol (5min) at room temperature and incubated with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 1h at room temperature to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody ab45977 at 5µg/ml and **ab72791** (Mouse monoclonal to alpha Tubulin - Loading Control) at 1ug/ml overnight at +4°C. The secondary antibodies used were **ab150081** used at 1 ug/ml (colored green) and **ab150116** (pseudo-colored red) used at 2ug/ml for 1 hour at room temperature. DAPI was used to stain the cell nuclei (colored blue) at a concentration of 1.43µM for 1 hour at room temperature.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Myeloperoxidase antibody (ab45977)

This image is courtesy of an Abreview submitted by Antibody Solutions Ltd.

ab45977 (1/50) staining Myeloperoxidase in paraffin-embedded human spleen tissue sections. Tissue underwent fixation in formaldehyde, heat-mediated antigen retrieval in citrate buffer pH 6.0 and blocking (5 minutes/peroxidase block and 10 minutes/protein block). For further experimental details please refer to abreview.



Western blot - Anti-Myeloperoxidase antibody
(ab45977)

Anti-Myeloperoxidase antibody (ab45977) at 1 µg/ml + HL60
(Human promyelocytic leukemia cell line) Whole Cell Lysate at 20
µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/50000 dilution

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 84 kDa

Observed band size: 84 kDa


Exposure time: 2 minutes

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 2% Bovine Serum Albumin before being incubated with ab45977 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution [ab133406](#)



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-Myeloperoxidase
antibody (ab45977)

IHC image of 45977 staining in human tonsil formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab45977, 1 µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western blot - Anti-Myeloperoxidase antibody (ab45977)

Anti-Myeloperoxidase antibody (ab45977) at 1 µg/ml + HL60 (Human promyelocytic leukemia cell line) Whole Cell Lysate at 20 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) preadsorbed (**ab97080**) at 1/5000 dilution

Predicted band size: 84 kDa

Observed band size: 84 kDa

Additional bands at: 12 kDa, 35 kDa, 75 kDa. We are unsure as to the identity of these extra bands.

Exposure time: 3 minutes

The mature myeloperoxidase protein is a tetramer of two heavy chains (60 kDa) and two light chains (12 kDa). Our immunogen sequence is within the myeloperoxidase light chain. In HL60 cells, ab45977 detects bands at approximately 84-kDa, corresponding to the expected MW of full-length Myeloperoxidase protein, and at 12-kDa, corresponding to the expected MW of myeloperoxidase light chain.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors