


Anti-N Cadherin antibody [8C11] ab19348

1 Abreviews **25 References** **画像数 3**

製品の概要

製品名	Anti-N Cadherin antibody [8C11]
製品の詳細	Mouse monoclonal [8C11] to N Cadherin
由来種	Mouse
アプリケーション	適用あり: ICC/IF, IHC-P
種交差性	交差種: Human, Bird 交差が予測される動物種: Rabbit, Hamster  非交差種: Mouse, Rat, Cow, Pig
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
エピトープ	The 8C11 monoclonal binds to the extracellular domain of N-cadherin between EC3 and EC4 (PubMed ID: 12604612).
ポジティブ・コントロール	IHC-P: Normal human heart tissue sections. ICC/IF: SH-SY5Y cells.
特記事項	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.02% Sodium azide Constituent: PBS
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	8C11

アイソタイプ	IgG1
軽鎖の種類	kappa

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab19348の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 5 µg/ml.
IHC-P		Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

ターゲット情報

機能	Cadherins are calcium dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. CDH2 may be involved in neuronal recognition mechanism. In hippocampal neurons, may regulate dendritic spine density.
配列類似性	Contains 5 cadherin domains.
細胞内局在	Cell membrane.

画像




IHC image of N cadherin staining in a section of formalin-fixed paraffin-embedded normal human heart performed on a Leica BOND™ system using the standard protocol F.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab19348, 1 µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with hematoxylin and mounted with DPX.

The inset secondary-only control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.




Immunocytochemistry/ Immunofluorescence - Anti-N Cadherin antibody [8C11] (ab19348)

ab19348 staining N-Cadherin in SH-SY5Y (Human neuroblastoma cell line from bone marrow) cells.

The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Tween for 5 mins and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab19348 at 5 µg/ml and **ab6046**, Rabbit polyclonal to beta Tubulin - Loading Control, at 1/1000 dilution. Cells were then incubated with **ab150117**, Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) at 1/1000 dilution (shown in green) and **ab150084**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 594) at 1/1000 dilution (shown in pseudocolor red). Nuclear DNA was labeled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunocytochemistry/ Immunofluorescence - Anti-N Cadherin antibody [8C11] (ab19348)

Immunofluorescence analysis of ARPE-19 (human retinal pigment epithelial) cells, staining N Cadherin (green) with ab19348, at 1/20 dilution.

ARPE-19 monolayer cultures were fixed in paraformaldehyde, permeabilized with 0.2% Triton X-100 for 15 min and blocked with 2% BSA for 30 min. Samples were incubated with primary antibody for 16 hours at 4°C before incubation with an Alexa Fluor® 488-conjugated donkey anti-mouse secondary IgG for 60 min.

Image from Chen HC et al., PLoS One. 2012;7(5):e36864. Epub 2012 May 9. Fig 1.; doi: 10.1371/journal.pone.0036864; May 9, 2012, PLoS One. 2012; 7(5): e36864. Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors