## abcam

#### Product datasheet

### Anti-N epsilon gamma glutamyl Lysine antibody [81D4] ab424

★★★★★ 2 Abreviews 17 References 画像数 2

#### 製品の概要

製品名 Anti-N epsilon gamma glutamyl Lysine antibody [81D4]

製品の詳細 Mouse monoclonal [81D4] to N epsilon gamma glutamyl Lysine

由来種 Mouse

アプリケーション 適用あり: IHC-P, ELISA, IHC-Fr

種交差性 交差種: Species independent

免疫原 Synthetic isopeptide.

特記事項 The ability of this antibody to fix complement is unknown. The isopeptide may be purchased from

Bachem Catalog number G-1970.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

#### 製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle.

רקלע"א Constituents: 2% Sucrose, 5.8% Sodium chloride, Phosphate Buffer

精製度 Protein A purified

**ポリ/モノ** モノクローナル

**クローン名** 81D4

ミエローマ unknown


アイソタイプ IgM

軽鎖の種類 kappa

# The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab424の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
IHC-P		1/50 - 1/800.
ELISA		Use at an assay dependent concentration.
IHC-Fr	<b>★★★★☆ (1)</b>	1/50 - 1/800.

#### 画像



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-N epsilon gamma glutamyl Lysine antibody [81D4] (ab424)

ab424 staining N epsilon gamma glutamyl Lysine in skin tissue sections.



Immunohistochemistry (Frozen sections) - Anti-N epsilon gamma glutamyl Lysine antibody [81D4] (ab424) ab424 staining N epsilon gamma glutamyl Lysine in kidney tissue sections by Immunohistochemistry (Frozen sections). Sections A through D were stained for insoluble tissue transglutaminase, while E through H were stained for N epsilon gamma glutamyl Lysine crosslink. A, B, E, F were control kidneys; C, D, G, H were diseased kidneys showing increased staining in the expanded interstitium, the mesangium and periglomerular.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors