

### Anti-Nanog antibody [NNG-811] ab62734

★★★★★ **3 Abreviews**   **40 References**   画像数 3

#### 製品の概要

製品名	Anti-Nanog antibody [NNG-811]
製品の詳細	Mouse monoclonal [NNG-811] to Nanog
由来種	Mouse
アプリケーション	<b>適用あり:</b> WB, ICC/IF, IP
種交差性	<b>交差種:</b> Human
免疫原	Recombinant full length human Nanog.
ポジティブ・コントロール	Extracts of NT2 cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	<p>pH: 7.40</p> <p>Preservative: 0.097% Sodium azide</p> <p>Constituent: 0.0268% PBS</p>
精製度	IgG fraction
特記事項(精製)	Purified Immunoglobulin
ポリ/モノ	モノクローナル
クローン名	NNG-811
アイソタイプ	IgG1


**The Abpromise guarantee** Abpromise保証は、 次のテスト済みアプリケーションにおけるab62734の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (1)	Use a concentration of 2 - 4 µg/ml. Detects a band of approximately 40 kDa (predicted molecular weight: 35 kDa).
ICC/IF		Use a concentration of 5 µg/ml.
IP		Use at an assay dependent concentration.


## ターゲット情報

機能	Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes (By similarity). Acts as a transcriptional activator or repressor (By similarity). Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3' (By similarity). When overexpressed, promotes cells to enter into S phase and proliferation.
組織特異性	Expressed in testicular carcinoma and derived germ cell tumors (at protein level). Expressed in fetal gonads, ovary and testis. Also expressed in ovary teratocarcinoma cell line and testicular embryonic carcinoma. Not expressed in many somatic organs and oocytes.
配列類似性	Belongs to the Nanog homeobox family. Contains 1 homeobox DNA-binding domain.
発生段階	Expressed in embryonic stem (ES) and carcinoma (EC) cells. Expressed in inner cell mass (ICM) of the blastocyst and gonocytes between 14 and 19 weeks of gestation (at protein level). Not expressed in oocytes, unfertilized oocytes, 2-16 cell embryos and early morula (at protein level). Expressed in embryonic stem cells (ES). Expression decreases with ES differentiation.
細胞内局在	Nucleus.

## 画像



3T3 mouse fibroblasts were transfected with a mammalian expression vector expressing human Nanog and stained with ab62734 (5 µg/mL) followed by Goat Anti-Mouse, Cy3 conjugate. A clear detection of Nanog can be seen in nuclei of transfected cells.



Western blot - Anti-Nanog antibody [NNG-811]  
(ab62734)

**Lane 1 :** Anti-Nanog antibody [NNG-811] (ab62734) at 10 µg/ml

**Lane 2 :** Anti-Nanog antibody [NNG-811] (ab62734) at 2 µg/ml

**Lane 3 :** Negative control – no primary antibody

**All lanes :** Cell extract of NT2 cells

Lysates/proteins at 20 µg per lane.


#### Secondary

**All lanes :** Goat Anti-Mouse IgG

Developed using the ECL technique.

**Predicted band size:** 35 kDa

**Observed band size:** 40 kDa



Immunoprecipitation - Anti-Nanog antibody [NNG-811] (ab62734)

Nanog was immunoprecipitated from NT2/D1 (human embryonal testis carcinoma cell line) whole cell extract using ab62734.

**Lane 1:** 5 µg ab62734 IP in NT2/D1 whole cell extract.

**Lane 2:** 10 µg ab62734 IP in NT2/D1 whole cell extract.

**Lane 3:** Negative control without cell extract.

**Lane 4:** NT2/D1 whole cell extract (input).

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors