abcam

Product datasheet

Anti-NAP1L1 antibody [2609C3a] ab58677

★★★★★★ 1 Abreviews 画像数 1

製品の概要

製品名 Anti-NAP1L1 antibody [2609C3a]

製品の詳細 Mouse monoclonal [2609C3a] to NAP1L1

由来種 Mouse

アプリケーション **適用あり**: WB

種交差性 交差種: Recombinant fragment **免疫原** Recombinant fragment (Human)

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

cycles.

バッファー pH: 7.40

Preservative: 0.05% Sodium azide

Constituents: PBS, 0.0225% Potassium chloride, 0.03% Potassium phosphate, 0.1312%

Sodium phosphate, 0.812% Sodium chloride, 1% BSA

精製度 Protein G purified

特記事項(精製) Purified using protein G column chromatography from culture supernatant of hybridoma cultured in

a medium containing bovine lgG-depleted (approximately 95%) fetal bovine serum.

ポリ/モノ モノクローナル

クローン名 2609C3a

アイソタイプ lgG1


The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab58677の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★ ☆☆ (1)	Use at an assay dependent concentration. Predicted molecular weight: 45 kDa.

ターゲット情報

機能	May be involved in modulating chromatin formation and contribute to regulation of cell proliferation.
組織特異性	Ubiquitously expressed.
配列類似性	Belongs to the nucleosome assembly protein (NAP) family.
ドメイン	The acidic domains are probably involved in the interaction with histones.
翻訳後修飾	Polyglutamylated by TTLL4, a modification that occurs exclusively on glutamate residues and results in polyglutamate chains on the gamma-carboxyl group. Some residues may also be monoglycylated but not polyglycylated due to the absence of functional TTLL10 in human.
細胞内局在	Nucleus. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

画像



Anti-NAP1L1 antibody [2609C3a] (ab58677)

Predicted band size: 45 kDa **Observed band size:** 38 kDa

Western blot analysis of immunised recombinant protein

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

• Replacement or refund for products not performing as stated on the datasheet

- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors