

### Anti-NCR1 antibody [MM0491-8F24] ab89877

★★★★★ [4 Abreviews](#) [1 References](#)

#### 製品の概要

製品名	Anti-NCR1 antibody [MM0491-8F24]
製品の詳細	Mouse monoclonal [MM0491-8F24] to NCR1
由来種	Mouse
アプリケーション	<b>適用あり:</b> IHC-Fr, Flow Cyt, WB
種交差性	<b>交差種:</b> Human
免疫原	Recombinant fragment corresponding to the extracellular domain of Human NCR1.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	Constituent: PBS
精製度	Protein G purified
特記事項 (精製)	The IgG fraction of culture supernatant was purified by Protein G affinity chromatography and sterile filtered through a 0.2 µm filter.
ポリ/モノ	モノクローナル
クローン名	MM0491-8F24
アイソタイプ	IgG2b

#### アプリケーション

The Abpromise guarantee

Abpromise保証は、 次のテスト済みアプリケーションにおけるab89877の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
IHC-Fr	★★★★★ (1)	Use at an assay dependent concentration.
Flow Cyt	★★★★★ (2)	Use at an assay dependent concentration. <b>ab170192</b> - Mouse monoclonal IgG2b, is suitable for use as an isotype control with this antibody.
WB	★★★★★ (1)	Use at an assay dependent concentration.

## ターゲット情報

機能	Cytotoxicity-activating receptor that may contribute to the increased efficiency of activated natural killer (NK) cells to mediate tumor cell lysis.
組織特異性	Selectively expressed by both resting and activated NK cells.
配列類似性	Belongs to the natural cytotoxicity receptor (NCR) family. Contains 2 Ig-like (immunoglobulin-like) domains.
翻訳後修飾	N-glycosylated. O-glycosylated.
細胞内局在	Cell membrane.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors