

Anti-NDUFS3 antibody [17D95] ab14711

★★★★★ **1 Abreviews** **98 References** 画像数 4

製品の概要

製品名	Anti-NDUFS3 antibody [17D95]
製品の詳細	Mouse monoclonal [17D95] to NDUFS3
由来種	Mouse
アプリケーション	適用あり: WB, Flow Cyt
種交差性	交差種: Mouse, Rat, Cow, Human
免疫原	Tissue, cells or virus. This information is considered to be commercially sensitive.
ポジティブ・コントロール	Human heart mitochondria. In Flow Cytometry, this antibody gave a positive signal in methanol fixed/Tween permeabilised HepG2 cells.
特記事項	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> <p>Product was previously marketed under the MitoSciences sub-brand.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 7.5 Preservative: 0.02% Sodium azide Constituent: HEPES buffered saline
精製度	Immunogen affinity purified
ポリ/モノ	モノクローナル
クローン名	17D95

アイソタイプ	IgG2a
軽鎖の種類	kappa

アプリケーション


The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab14711の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (1)	Use a concentration of 0.5 - 1 µg/ml. Detects a band of approximately 26 kDa (predicted molecular weight: 30 kDa).
Flow Cyt		Use 1µg for 10 ⁶ cells. ab170191 - Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody.

ターゲット情報

機能	Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.
配列類似性	Belongs to the complex I 30 kDa subunit family.
細胞内局在	Mitochondrion inner membrane.

画像



All lanes : Anti-NDUFS3 antibody [17D95] (ab14711) at 1 µg/ml

- Lane 1 :** Human brain tissue lysate - total protein ([ab29466](#))
- Lane 2 :** Human heart tissue lysate - total protein ([ab29431](#))
- Lane 3 :** Heart (Mouse) Tissue Lysate
- Lane 4 :** HepG2 (Human hepatocellular liver carcinoma cell line) Whole Cell Lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Mouse IgG H&L (HRP) preadsorbed ([ab97040](#)) at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.


Predicted band size: 30 kDa

Observed band size: 26 kDa

Additional bands at: 57 kDa. We are unsure as to the identity of these extra bands.

Exposure time: 1 minute

The band observed at 26 kDa could potentially be a cleaved form of NDUFS3 due to the presence of a 36 amino acid transit peptide.



Western blot - Anti-NDUFS3 antibody [17D95] (ab14711)

Image is courtesy of an anonymous AbReview

All lanes : Anti-NDUFS3 antibody [17D95] (ab14711) at 1/1000 dilution

Lanes 1-2 : Whole cell lysates from NDUFS3 KO cells.

Lanes 3-4 : Whole cell lysates from NDUFS3 WT cells.

Lanes 5-6 : Isolated mitochondria from NDUFS3 KO cells.

Lanes 7-8 : Isolated mitochondria from NDUFS3 WT cells.

Lysates/proteins at 50 µg per lane.

Secondary

All lanes : Goat anti-mouse polyclonal HRP conjugate at 1/5000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 30 kDa


Additional bands at: 22 kDa (possible non-specific binding), 40 kDa (possible non-specific binding)

Exposure time: 1 minute



Flow Cytometry - Anti-NDUFS3 antibody [17D95]
(ab14711)

Overlay histogram showing HepG2 cells stained with ab14711 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab14711, 1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2a [ICIGG2A] ([ab91361](#), 1µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed.



Western blot - Anti-NDUFS3 antibody [17D95]
(ab14711)

All lanes : Anti-NDUFS3 antibody [17D95] (ab14711)

Lane 1 : Isolated mitochondria from Human heart at 5 µg

Lane 2 : Isolated mitochondria from Bovine heart at 4 µg

Lane 3 : Isolated mitochondria from Rat heart at 10 µg

Lane 4 : Isolated mitochondria from Mouse heart at 10 µg

Secondary

All lanes : Goat anti-Mouse IgG

Predicted band size: 30 kDa

Observed band size: 26 kDa

Extra bands in the mouse sample (lane 4) are due to the reaction of the IgG-specific goat anti-mouse secondary antibody with residual mouse blood in the heart tissue, as it is very difficult to entirely remove the blood from these small organs.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors