

Anti-Neisseria gonorrhoeae antibody ab19962

2 References

製品の概要

製品名	Anti-Neisseria gonorrhoeae antibody
製品の詳細	Rabbit polyclonal to Neisseria gonorrhoeae
由来種	Rabbit
特異性	Reacts to all antigens of Neisseria gonorrhoeae. Antiserum is not absorbed and may react with related microorganisms.
アプリケーション	適用あり: ICC/IF
種交差性	交差種: Neisseria gonorrhoeae
免疫原	Tissue, cells or virus corresponding to Human Neisseria gonorrhoeae. Whole Neisseria gonorrhoeae.ATCC 31426
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.20</p> <p>Preservative: 0.1% Sodium azide</p> <p>Constituent: 0.0268% PBS</p>
精製度	Protein A purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab19962の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use at an assay dependent concentration.

ターゲット情報

関連性	Neisseria gonorrhoeae infections are acquired by sexual contact and usually affect the mucous membranes of the urethra in males and the endocervix and urethra in females, although the infection may disseminate to a variety of tissues. The pathogenic mechanism involves the attachment of the bacterium to nonciliated epithelial cells via pili (fimbriae) and the production of lipopolysaccharide endotoxin. Similarly, the lipopolysaccharide of Neisseria meningitidis is highly toxic, as it has an additional virulence factor in the form of its antiphagocytic capsule. Both pathogens produce IgA proteases which promote virulence. Many normal individuals may harbor Neisseria meningitidis in the upper respiratory tract, but Neisseria gonorrhoeae is never part of the normal flora and is only found after sexual contact with an infected person (or direct contact, in the case of infections in the newborn).
-----	---

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors