abcam

Product datasheet

Anti-Nuclear Matrix Protein p84 antibody [EPR5661] ab125019

יעלאעבע RabMAb

1 References 画像数4

製品の概要

製品名 Anti-Nuclear Matrix Protein p84 antibody [EPR5661]

製品の詳細 Rabbit monoclonal [EPR5661] to Nuclear Matrix Protein p84

由来種 Rabbit

アプリケーション 適用あり: WB, IHC-P, ICC/IF

適用なし: Flow Cvt or IP

種交差性 交差種: Human

免疫原 Synthetic peptide within Human Nuclear Matrix Protein p84 aa 1-100. The exact sequence is

proprietary.

ポジティブ・コントロール HeLa, U-87 MG and 293T cell lysates; Human stomach adenocarcinoma tissue; 293T cells

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with

these species. Please contact us for more information.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.

バッファー pH: 7.20

Preservative: 0.01% Sodium azide

Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture

supernatant

精製度 Protein A purified

ポリ/モノ モノクローナル **ウローン名** EPR5661

アイソタイプ IgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab125019の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/1000 - 1/10000. Detects a band of approximately 84 kDa (predicted molecular weight: 76 kDa).
IHC-P		1/50 - 1/100. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
ICC/IF		1/100 - 1/250.

追加情報

Is unsuitable for Flow Cyt or IP.

ターゲット情報

組織特異性

機能

Component of the THO subcomplex of the TREX complex. The TREX complex specifically associates with spliced mRNA and not with unspliced pre-mRNA. It is recruited to spliced mRNAs by a transcription-independent mechanism. Binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export. The recruitment occurs via an interaction between THOC4 and the cap-binding protein NCBP1. DDX39B functions as a bridge between THOC4 and the THO complex. The TREX complex is essential for the export of Kaposi's sarcoma-associated herpesvirus (KSHV) intronless mRNAs and infectious virus production. The recruitment of the TREX complex to the intronless viral mRNA occurs via an interaction between KSHV ORF57 protein and THOC4.

Regulates transcriptional elongation of a subset of genes. Participates in an apoptotic pathway which is characterized by activation of caspase-6, increases in the expression of BAK1 and BCL2L1 and activation of NF-kappa-B. This pathway does not require p53/TP53, nor does the presence of p53/TP53 affect the efficiency of cell killing. Activates a G2/M cell cycle checkpoint prior to the onset of apoptosis. Apoptosis is inhibited by association with RB1.

Ubiquitous. Expressed in various cancer cell lines. Expressed at very low levels in normal breast epithelial cells and highly expressed in breast tumors. Expression is strongly associated with an aggressive phenotype of breast tumors and expression correlates with tumor size and the metastatic state of the tumor progression.

配列類似性 Contains 1 death domain.


ドメイン An intact death domain is needed for apoptosis.

翻訳後修飾 Expression is altered specifically during apoptosis and is accompanied by the appearance of

novel forms with smaller apparent molecular mass.

細胞内局在Cytoplasm and Nucleus speckle. Nucleus > nucleoplasm. Nucleus matrix. Cytoplasm. Can shuttle between the nucleus and cytoplasm. Nuclear localization is required for induction of apoptotic cell

画像



Western blot - Anti-Nuclear Matrix Protein p84 antibody [EPR5661] (ab125019)

All lanes : Anti-Nuclear Matrix Protein p84 antibody [EPR5661] (ab125019) at 1/1000 dilution

Lane 1: Hela cell lysate

Lane 2 : U-87 MG (Human glioblastoma-astrocytoma epithelial cell line) cell lysate


Lane 3: 293T (Human embryonic kidney epithelial cell) cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes: HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 76 kDa **Observed band size:** 84 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Nuclear Matrix Protein p84 antibody [EPR5661] (ab125019)

ab125019, at 1/50 dilution, staining Nuclear Matrix Protein p84 in paraffin-embedded Human stomach adenocarcinoma tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Immunocytochemistry/ Immunofluorescence - Anti-Nuclear Matrix Protein p84 antibody [EPR5661] (ab125019)

ab125019, at 1/100 dilution, staining Nuclear Matrix Protein p84 in 293T cells by Immunofluorescence.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

Guarantee only valid for products bought direct from Abcam or one of our authorized distributors