

# Anti-Nuclear Receptor Corepressor NCoR antibody ab58396

**5 References** [画像数 2](#)

### 製品の概要

製品名	Anti-Nuclear Receptor Corepressor NCoR antibody
製品の詳細	Rabbit polyclonal to Nuclear Receptor Corepressor NCoR
由来種	Rabbit
特異性	This antibody detects endogenous levels of total NCoR1 protein.
アプリケーション	<b>適用あり:</b> WB, IHC-P
種交差性	<b>交差種:</b> Human
免疫原	Synthetic peptide derived from the internal sequence of Nuclear Receptor Corepressor NCoR (Human)
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	<p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituents: PBS, 50% Glycerol (glycerin, glycerine), 0.87% Sodium chloride</p>
精製度	Without Mg+2 and Ca+2
特記事項 (精製)	Immunogen affinity purified
ポリ/モノ	Purified from rabbit antiserum.
アイソタイプ	ポリクローナル
	IgG

## アプリケーション


**The Abpromise guarantee** Abpromise保証は、 次のテスト済みアプリケーションにおけるab58396の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/500 - 1/1000. Predicted molecular weight: 270 kDa.
IHC-P		1/50 - 1/100.

## ターゲット情報

機能	Mediates transcriptional repression by certain nuclear receptors. Part of a complex which promotes histone deacetylation and the formation of repressive chromatin structures which may impede the access of basal transcription factors.
配列類似性	Belongs to the N-CoR nuclear receptor corepressors family. Contains 2 SANT domains.
ドメイン	The N-terminal region contains three independent domains that are capable of mediating transcriptional repression (RD1, RD2 and RD3). The C-terminal region contains two separate nuclear receptor-interacting domains (ID1 and ID2), each of which contains a conserved sequence referred to as the CORNR box. This motif is necessary and sufficient for binding to unligated nuclear hormone receptors, while sequences flanking the CORNR box determine the precise nuclear hormone receptor specificity.
翻訳後修飾	Ubiquitinated; mediated by SIAH2 and leading to its subsequent proteasomal degradation.
細胞内局在	Nucleus.

## 画像




**All lanes :** Anti-Nuclear Receptor Corepressor NCoR antibody (ab58396) at 1/500 dilution

**Lane 1 :** MDA-MB-435 cell extract with no immunizing peptide

**Lane 2 :** MDA-MB-435 cell extract with immunizing peptide

**Predicted band size:** 270 kDa



ab58396, at 1/50 dilution, staining human Nuclear Receptor Corepressor NCoR in breast carcinoma tissue by immunohistochemistry in the presence or absence of the immunizing peptide. Paraffin embedded tissue.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors