

# Anti-NUP98 antibody [2H10] - Nuclear Pore Marker ab50610

★★★★☆ **5 Abreviews** **25 References** **画像数 5**

### 製品の概要

---

製品名	Anti-NUP98 antibody [2H10] - Nuclear Pore Marker
製品の詳細	Rat monoclonal [2H10] to NUP98 - Nuclear Pore Marker
由来種	Rat
アプリケーション	<b>適用あり:</b> WB, ICC/IF
種交差性	<b>交差種:</b> Mouse, Rat, Human, African green monkey
免疫原	Recombinant fragment corresponding to Human NUP98 aa 1-466.
ポジティブ・コントロール	WB: HeLa nuclear lysate. Jurkat, HeLa, COS-7, NIH/3T3 and SH-SY5Y cell lysate. ICC/IF: HeLa and NIH/3T3 cells.
特記事項	<p>This product was changed from ascites to tissue culture supernatant on 17 May 2019. Please note that the dilutions may need to be adjusted accordingly. If you have any questions, please do not hesitate to contact our scientific support team.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

---

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.097% Sodium azide Constituent: 0.0268% PBS
精製度	Tissue culture supernatant
特記事項(精製)	Purified from TCS.
ポリ/モノ	モノクローナル

クローン名	2H10
ミエローマ	Sp2
アイソタイプ	IgG2c
軽鎖の種類	kappa


## アプリケーション

**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab50610の使用に適用されます  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (2)	1/1000. Detects a band of approximately 98 kDa.
ICC/IF	★★★★★ (1)	Use at an assay dependent concentration. Customers have reported that Paraformaldehyde/Triton x-100 fixation provides better results, with sharp, regularly punctuate perinuclear signals. In MetOH fixed cells, the signal intensity can be somewhat lower and fuzzier and that single nucleoporin dots can be harder to distinguish around nuclear chromatin. Please see images below.

## ターゲット情報

<b>機能</b>	Nup98 and Nup96 play a role in the bidirectional transport across the nucleoporin complex (NPC). The repeat domain in Nup98 has a direct role in the transport.
<b>関連疾患</b>	Note=A chromosomal aberration involving NUP98 is found in a form of acute myeloid leukemia. Translocation t(7;11)(p15;p15) with HOXA9. Translocation t(11;17)(p15;p13) with PHF23. Note=A chromosomal aberration involving NUP98 is found in childhood acute myeloid leukemia. Translocation t(5;11)(q35;p15.5) with NSD1. Translocation t(8;11)(p11.2;p15) with WHSC1L1. Note=A chromosomal aberration involving NUP98 is found in a form of therapy-related myelodysplastic syndrome. Translocation t(11;20)(p15;q11) with TOP1. Note=A chromosomal aberration involving NUP98 is found in a form of T-cell acute lymphoblastic leukemia (T-ALL). Translocation t(3;11)(q12.2;p15.4) with LNP1. Note=A chromosomal aberration involving NUP98 is associated with pediatric acute myeloid leukemia (AML) with intermediate characteristics between M2-M3 French-American-British (FAB) subtypes. Translocation t(9;11)(p22;p15) with PSIP1/LEDGF. The chimeric transcript is an in-frame fusion of NUP98 exon 8 to PSIP1/LEDGF exon 4.
<b>配列類似性</b>	Belongs to the nucleoporin GLFG family. Contains 1 peptidase S59 domain.
<b>ドメイン</b>	Contains G-L-F-G repeats.
<b>翻訳後修飾</b>	Isoform 1 to isoform 4 are autoproteolytically cleaved to yield Nup98 and Nup96 or Nup98 only, respectively. Cleaved Nup98 is necessary for the targeting of Nup98 to the nuclear pore and the interaction with Nup96.
<b>細胞内局在</b>	Nucleus > nuclear pore complex. Nucleus membrane. Nup96 is localized to the nucleoplasmic side of the nuclear pore complex, at or near the nucleoplasmic basket.



Immunocytochemistry/ Immunofluorescence - Anti-NUP98 antibody [2H10] - Nuclear Pore Marker (ab50610)


Di Nunzio et al PLoS One. 2012;7(9):e46037. doi: 10.1371/journal.pone.0046037. Epub 2012 Sep 25. Fig 1. Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>

**Lentiviral vector-encoded shRNAs achieve efficient knock-down of human nucleoporins and have negligible cytotoxic or cytostatic effects.**

HeLa cells ( $4 \times 10^6$ ) were transduced with lentiviral vectors (MOI 50) encoding shRNAs specific for the indicated nucleoporins and used at 2 days p.t for Nup153 shRNA and 5 days p.t for all others.

**(Panel B)** Subcellular localisation of nuclear pore components upon nucleoporin knock-down was tested by confocal fluorescence microscopy of LV- (Control) and LV-shRNA transduced cells using specific anti-Nup antibodies. Images were acquired on the same day with the same conditions and are representative of two independent experiments.

This image was generated using the ascites version of the product.



Western blot - Anti-NUP98 antibody [2H10] - Nuclear Pore Marker (ab50610)

**All lanes :** Anti-NUP98 antibody [2H10] - Nuclear Pore Marker (ab50610) at 1 µg/ml

**Lane 1 :** HeLa (human epithelial cell line from cervix adenocarcinoma) nuclear lysate

**Lane 2 :** HeLa (human epithelial cell line from cervix adenocarcinoma) cell lysate

**Lane 3 :** Jurkat (human T cell leukemia cell line from peripheral blood) cell lysate

**Lane 4 :** SH-SY5Y (human neuroblastoma cell line from bone marrow) cell lysate

**Lane 5 :** COS-7 (african green monkey kidney fibroblast-like cell line) cell lysate

**Lane 6 :** NIH/3T3 (mouse embryo fibroblast cell line) cell lysate

**Lane 7 :** P19 cell lysate


**Lane 8 :** NRK cell lysate

**Secondary**

**All lanes :** Goat Anti-Mouse IgG-Peroxidase

Developed using the ECL technique.

This image was generated using the ascites version of the product.




ab50610 (1/100) staining NUP98 in HeLa (Human epithelial cell line from cervix adenocarcinoma) cells (green).

Cells were fixed with paraformaldehyde/Triton X-100 [10 min in PTEMF buffer (20mM PIPES, 1mM MgCl<sub>2</sub>, 10mM EGTA, 4% PFA) /0.2% Triton X-100 at room temperature] or methanol (6 min in methanol -20 °C , followed by 3 washes in 1x PBS) and counterstained with DAPI in order to highlight the nucleus (blue).

This image was generated using the ascites version of the product.

Immunocytochemistry/ Immunofluorescence - Anti-NUP98 antibody [2H10] - Nuclear Pore Marker (ab50610)

Image and protocol courtesy of Rosamaria Mangiacasale, Marilena Ciciarello and Patrizia Lavia, Univ Rome La Sapienza, Italy




ab50610 (1/100) staining NUP98 in NIH/3T3 (Mouse embryo fibroblast cell line) cells (green).

Cells were fixed with paraformaldehyde/Triton X-100 (10 min in PTEMF buffer (20mM PIPES, 1mM MgCl<sub>2</sub>, 10mM EGTA, 4% PFA) /0.2% Triton X-100 at room temperature) or methanol (6 min in methanol -20 °C , followed by 3 washes in 1x PBS) and counterstained with DAPI in order to highlight the nucleus (blue).

This image was generated using the ascites version of the product.

Immunocytochemistry/ Immunofluorescence - Anti-NUP98 antibody [2H10] - Nuclear Pore Marker (ab50610)

Image and protocol courtesy of Rosamaria Mangiacasale, Marilena Ciciarello and Patrizia Lavia, Univ Rome La Sapienza, Italy



Immunocytochemistry/ Immunofluorescence - Anti-NUP98 antibody [2H10] - Nuclear Pore Marker (ab50610)

This image is courtesy of an anonymous Abreview.

Paraformaldehyde-fixed, 0.5% Triton X-100 permeabilized HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) cells stained for NUP98 (green) using ab50610 at 1/200 dilution in ICC/IF, followed by Donkey Anti-Rat Alexa Fluor<sup>®</sup> 488.

This image was generated using the ascites version of the product.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors