abcam

Product datasheet

Anti-Oct-2 antibody [EP543Y] ab33919

RabMAb

3 References 画像数 1

製品の概要

製品名 Anti-Oct-2 antibody [EP543Y]

製品の詳細 Rabbit monoclonal [EP543Y] to Oct-2

由来種 Rabbit

特異性 This antibody may detect the isoform 2.

アプリケーション 適用あり: WB

適用なし: ChIP or Flow Cyt

種交差性 交差種: Human

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール Raji cell lysate

特記事項 Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit

monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with

these species. Please contact us for more information.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle.

バッファー pH: 7.20

Preservative: 0.01% Sodium azide

Constituents: 49% PBS, 50% Glycerol (glycerin, glycerine), 0.05% BSA

1

精製度 Protein A purified

ポリ/モノ モノクローナル

クローン名 EP543Y

アイソタイプ lgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab33919の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/20000 - 1/50000. Predicted molecular weight: 51 kDa.

追加情報 Is unsuitable for ChIP or Flow Cyt.

ターゲット情報

機能 Transcription factor that specifically binds to the octamer motif (5'-ATTTGCAT-3'). Regulates

transcription in a number of tissues in addition to activating immunoglobulin gene expression.

Modulates transcription transactivation by NR3C1, AR and PGR. Isoform 5 activates the U2 small

nuclear RNA (snRNA) promoter.

組織特異性 Isoform 3 is B-cell specific. Isoform 5 is expressed in B-cells and the immunoglobulin-expressing

T-cell line MOLT-4, but not in the T-cell line BW5147.


配列類似性 Belongs to the POU transcription factor family. Class-2 subfamily.

Contains 1 homeobox DNA-binding domain.

Contains 1 POU-specific domain.

細胞内局在 Cytoplasm. Nucleus.

画像



Anti-Oct-2 antibody [EP543Y] (ab33919) at 1/50000 dilution + Raji cell lysate

Predicted band size: 51 kDa **Observed band size:** 55 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors