

Anti-Otoferlin antibody [13A9] ab53233

★★★★★ [3 Abreviews](#) [33 References](#) [画像数 2](#)

製品の概要

製品名	Anti-Otoferlin antibody [13A9]
製品の詳細	Mouse monoclonal [13A9] to Otoferlin
由来種	Mouse
特異性	This antibody reacts specifically with human Otoferlin protein (220 kDa).
アプリケーション	適用あり: Flow Cyt, ICC/IF
種交差性	交差種: Mouse, Human
免疫原	Fusion protein corresponding to Human Otoferlin aa 1-400. Database link: Q9HC10
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Constituents: 1.21% Tris, 0.75% Glycine, 2% Sucrose
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	13A9
アイソタイプ	IgG1
軽鎖の種類	kappa

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab53233の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 1µg for 10 ⁶ cells. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
ICC/IF	★★★★★ (1)	1/100 - 1/500.

ターゲット情報


機能	Key calcium ion sensor involved in the Ca(2+)-triggered synaptic vesicle-plasma membrane fusion and in the control of neurotransmitter release at these output synapses. Interacts in a calcium-dependent manner to the presynaptic SNARE proteins at ribbon synapses of cochlear inner hair cells (IHCs) to trigger exocytosis of neurotransmitter. Also essential to synaptic exocytosis in immature outer hair cells (OHCs). May also play a role within the recycling of endosomes.
組織特異性	Isoform 1 and isoform 3 are found in adult brain. Isoform 2 is expressed in the fetus and in adult brain, heart, placenta, skeletal muscle and kidney.
関連疾患	Defects in OTOF are the cause of deafness autosomal recessive type 9 (DFNB9) [MIM:601071]. DFNB9 is a form of sensorineural hearing loss. Sensorineural deafness results from damage to the neural receptors of the inner ear, the nerve pathways to the brain, or the area of the brain that receives sound information. Defects in OTOF are a cause of non-syndromic auditory neuropathy autosomal recessive (NSRAN) [MIM:601071]. NSRAN is a form of sensorineural hearing impairment with absent or severely abnormal auditory brainstem response but normal otoacoustic emissions. Auditory neuropathies result from a lesion in the area including the inner hair cells, connections between the inner hair cells and the cochlear branch of the auditory nerve, the auditory nerve itself and auditory pathways of the brainstem. In some cases NSRAN phenotype can be temperature sensitive.
配列類似性	Belongs to the ferlin family. Contains 4 C2 domains.
細胞内局在	Cytoplasmic vesicle > secretory vesicle > synaptic vesicle membrane. Basolateral cell membrane. Endoplasmic reticulum membrane. Cell membrane. Detected at basolateral cell membrane with synaptic vesicles surrounding the ribbon and at the presynaptic plasma membrane in the inner hair cells (IHCs). Colocalizes with GPR25 and RAB8B in inner hair cells.

画像



Flow Cytometry - Anti-Otoferlin antibody [13A9] (ab53233)

Overlay histogram showing SHSY-5Y cells stained with ab53233 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab53233, 1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] ([ab91353](#), 2µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in SHSY-5Y cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Immunocytochemistry/ Immunofluorescence - Anti-Otoferlin antibody [13A9] (ab53233)

ICC/IF analysis of mouse inner ear tissue labeling Otoferlin with [ab181781](#). At 30 days postnatal, Otoferlin is detected in the hair cells of the crista ampullaris (CA), utricular macula (UM) and saccular macula (SM).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors