

Anti-p63 Protein + HMW Cytokeratin antibody [4A4, 34] ab58514

製品の概要

| | |
|--------------|---|
| 製品名 | Anti-p63 Protein + HMW Cytokeratin antibody [4A4, 34] |
| 製品の詳細 | Mouse monoclonal [4A4, 34] to p63 Protein + HMW Cytokeratin |
| 由来種 | Mouse |
| アプリケーション | 適用あり: IHC-P |
| 種交差性 | 交差種: Human |
| 免疫原 | Tissue, cells or virus corresponding to Human p63 Protein + HMW Cytokeratin. |
| ポジティブ・コントロール | Normal prostate tissue or skin |
| 特記事項 | <p>ab58514 isotypes: IgG2a + IgG1</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

製品の特性

| | |
|--------|---|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle. |
| バッファー | <p>pH: 7.30</p> <p>Preservative: 0.09% Sodium azide</p> <p>Constituent: PBS</p> |
| ポリ/モノ | モノクローナル |
| クローン名 | 4A4, 34 |
| アイソタイプ | IgG1 |

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab58514の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|------|
| IHC-P | | |

追加情報

IHC-P: ready to use, 100 µl working solution / test. Antigen retrieval requires unmasking, TEC buffer (Tris/EDTA/Citrate) pH 8 in a pressure cooker at 100°C 20-40 minutes

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

関連性

The p63 and high molecular weight cytokeratin cocktail is a marker for basal cells. The p63 protein, a homologue of the tumour-suppressor p53, is highly expressed in the nuclei of basal or progenitor layer of many epithelial tissues. p63 is detected in prostate basal cells in normal prostate glands and PIN. Other markers for basal cells are high molecular cytokeratins (CK 1, 5, 10 and 14). The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells.

細胞内局在

Cytoplasmic and Nuclear

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors