


Anti-PAPD5 antibody ab118341

★★★★★ [1 Abreviews](#) [画像数 1](#)

製品の概要

製品名	Anti-PAPD5 antibody
製品の詳細	Goat polyclonal to PAPD5
由来種	Goat
特異性	ab118341 is expected to recognize both reported isoforms (NP_001035374.2; NP_001035375.2).
アプリケーション	適用あり: WB
種交差性	交差種: Mouse 交差が予測される動物種: Horse, Cow, Dog, Human 
免疫原	Synthetic peptide corresponding to Human PAPD5 aa 450-550 (internal sequence) (Cysteine residue). NP_001035374.2; NP_001035375.2 Run BLAST with ExPASy Run BLAST with NCBI
ポジティブ・コントロール	WB: Mouse and rat spleen lysate.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 0.5% BSA, Tris buffered saline
精製度	Immunogen affinity purified
特記事項 (精製)	ab118341 is purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

ポリモノ
アイソタイプ

ポリクローナル
IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab118341の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB	★☆☆☆☆ (1)	Use a concentration of 0.2 - 0.6 µg/ml. Detects a band of approximately 63 kDa (predicted molecular weight: 63 kDa). 1 hour primary incubation is recommended for this product.

ターゲット情報

関連性 Plays a role in replication-dependent histone mRNA degradation. May be involved in the terminal uridylation of mature histone mRNAs before their degradation is initiated. DNA polymerase, probably involved in DNA repair. May play a role in sister chromatid cohesion

細胞内局在 Nuclear

画像



Anti-PAPD5 antibody (ab118341) at 0.2 µg/ml (for 1 hour) + Mouse spleen lysate (in RIPA buffer) at 35 µg

Developed using the ECL technique.

Predicted band size: 63 kDa

Observed band size: 70 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors