

### Anti-PARP1 antibody [E102] ab32138

KO 評価済 リコンビナント RabMAb

★★★★☆ 6 Abreviews 116 References 画像数 8

#### 製品の概要

製品名	Anti-PARP1 antibody [E102]
製品の詳細	Rabbit monoclonal [E102] to PARP1
由来種	Rabbit
特異性	This antibody recognises both pro-form and p25 cleaved form of PARP1.
アプリケーション	<b>適用あり:</b> WB, IHC-P, Flow Cyt (Intra), ICC/IF <b>適用なし:</b> IP
種交差性	<b>交差種:</b> Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Wild type HAP1 whole cell lysate; HeLa whole cell lysate ( <b>ab150035</b> ); HEK-293T cell lysate; IHC-P: Human breast carcinoma tissue; ICC/IF: HeLa cells; Flow Cyt (intra): HeLa cells, HAP1 cells. WB: Jurkat whole cell lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide

	Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	E102
アイソタイプ	IgG

## アプリケーション

**The Abpromise guarantee** **Abpromise保証は、次のテスト済みアプリケーションにおけるab32138の使用に適用されます**  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB	★★★★★ (4)	1/1000 - 1/10000. Predicted molecular weight: 113 kDa. Existing as a 113 kDa nuclear protein, PARP1 is cleaved between amino acids Asp214 and Gly215 to yield two fragments of 29 kDa (N-terminal catalytic domain) and 85 kDa (C-terminal DNA-binding domain)
IHC-P	★★★★★ (1)	1/200. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. See <b><u>IHC antigen retrieval protocols.</u></b>  <b>For unpurified use at 1/25.</b>
Flow Cyt (Intra)		1/20 - 1/50. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF	★★★★★ (1)	1/100.

**追加情報** Is unsuitable for IP.

## ターゲット情報

機能	Involved in the base excision repair (BER) pathway, by catalyzing the poly(ADP-ribosyl)ation of a limited number of acceptor proteins involved in chromatin architecture and in DNA metabolism. This modification follows DNA damages and appears as an obligatory step in a detection/signaling pathway leading to the reparation of DNA strand breaks. Mediates the poly(ADP-ribosyl)ation of APLF and CHFR. Positively regulates the transcription of MTUS1 and negatively regulates the transcription of MTUS2/TIP150.
配列類似性	Contains 1 BRCT domain. Contains 1 PARP alpha-helical domain. Contains 1 PARP catalytic domain. Contains 2 PARP-type zinc fingers.
翻訳後修飾	Phosphorylated by PRKDC. Phosphorylated upon DNA damage, probably by ATM or ATR. Poly-ADP-ribosylated by PARP2. Poly-ADP-ribosylation mediates the recruitment of CHD1L to DNA damage sites. S-nitrosylated, leading to inhibit transcription regulation activity.

画像




Flow cytometry overlay histogram showing wild-type Hap1 (green line) and PARP1 knockout Hap1 stained with ab32138 (red line). The cells were fixed with 4% formaldehyde (10 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10% normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab32138) ( $1 \times 10^6$  in 100 $\mu$ l at 0.04  $\mu$ g/ml (1/55750)) for 30min at 22°C.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min at 22°C

Isotype control antibody Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control was used at the same concentration and conditions as the primary antibody (wild-type Hap1 - black line, PARP1 knockout Hap1 - grey line). Unlabelled sample was also used as a control (this line is not shown for the purpose of simplicity).

Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter.



**All lanes :** Anti-PARP1 antibody [E102] (ab32138) at 1/1000 dilution (Purified)

**Lane 1 :** Untreated Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate


**Lane 2 :** Jurkat (Human T cell leukemia T lymphocyte) treated with 1 $\mu$ M staurosporine for 4 hours whole cell lysate

**Secondary**

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution


**Predicted band size:** 113 kDa

pro-form: 116kDa; p25 caspases cleaved form: 25kDa; proteolysis cleaved fragments: 58kDa and 42kDa



Immunocytochemistry/ Immunofluorescence - Anti-PARP1 antibody [E102] (ab32138)

Immunocytochemistry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling PARP1 with purified ab32138 at 1/100 dilution (1.0 µg/mL). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% TritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.5 µg/mL). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody at 1/1000 (2 µg/mL) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Western blot - Anti-PARP1 antibody [E102] (ab32138)

**Lane 1:** Wild type HAP1 whole cell lysate (20 µg)


**Lane 2:** PARP1 knockout HAP1 whole cell lysate (20 µg)

**Lane 3:** HeLa whole cell lysate (20 µg)

**Lane 4:** MCF7 whole cell lysate (20 µg)


**Lanes 1 - 4:** Merged signal (red and green). Green - ab32138 observed at 125 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

ab32138 was shown to specifically react with PARP1 when PARP1 knockout samples were used. Wild-type and PARP1 knockout samples were subjected to SDS-PAGE. ab32138 and [ab8245](#) (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/10 000 dilution respectively. Blots were developed with 800CW Goat anti Rabbit and 680CW Goat anti Mouse secondary antibodies at 1/10 000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PARP1 antibody [E102] (ab32138)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue sections labeling PARP1 with purified ab32138 at 1/200 dilution (0.51 µg/mL). Heat mediated antigen retrieval was performed using Perform heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1/0 dilution. PBS instead of the primary antibody was used as the negative control.



Western blot - Anti-PARP1 antibody [E102] (ab32138)

**All lanes :** Anti-PARP1 antibody [E102] (ab32138) at 1/1000 dilution

**Lane 1 :** Wild-type HEK-293T cell lysate

**Lane 2 :** PARP1 knockout HEK-293T cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


**Predicted band size:** 113 kDa

**Observed band size:** 113 kDa

**Lanes 1- 2:** Merged signal (red and green). Green - ab32138 observed at 113 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) observed at 37 kDa.





ab32138 was shown to react with PARP1 in wild-type HEK-293T cells in western blot. Loss of signal was observed when knockout cell line **ab266598** (knockout cell lysate **ab257017**) was used. Wild-type HEK-293T and PARP1 knockout HEK-293T cell lysates were subjected to SDS-PAGE. ab32138 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (**ab216776**) secondary antibodies at

1 in 20000 dilution for 1 hour at room temperature before imaging.



Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labelling PARP1 with purified ab32138 at 1/20 dilution (10 µg/mL) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabelled control - Cell without incubation with primary antibody and secondary antibody (Blue).

Why choose a recombinant antibody?

 <p><b>Research with confidence</b> Consistent and reproducible results</p>	 <p><b>Long-term and scalable supply</b> Recombinant technology</p>
 <p><b>Success from the first experiment</b> Confirmed specificity</p>	 <p><b>Ethical standards compliant</b> Animal-free production</p>

Anti-PARP1 antibody [E102] (ab32138)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors