

Anti-PAX8 antibody ab97477

★★★★★ [2 Abreviews](#) [6 References](#) [画像数 8](#)

製品の概要

製品名	Anti-PAX8 antibody
製品の詳細	Rabbit polyclonal to PAX8
由来種	Rabbit
アプリケーション	適用あり: WB, ICC/IF, ChIP, IHC-P, IP
種交差性	交差種: Mouse, Rat, Human
免疫原	Recombinant protein fragment corresponding to a region within amino acids 1 and 214 of PAX8 (Uniprot: Q06710).
ポジティブ・コントロール	ChIP: HEK-293T cells; WB: HEK-293T whole cell lysate; IHC-P: DLD1 xenograft, mouse liver and rat thyroid gland tissue; IF: A549 whole cells. IP: HEK-293T whole cell lysate.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.00</p> <p>Preservative: 0.025% Proclin 300</p> <p>Constituents: 79% PBS, 20% Glycerol (glycerin, glycerine)</p>
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab97477の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (1)	1/500 - 1/2000. Predicted molecular weight: 48 kDa.
ICC/IF	★★★★★ (1)	1/100 - 1/1000.
ChIP		Use at an assay dependent concentration.
IHC-P		1/100 - 1/1000.
IP		1/500 - 1/1000.

ターゲット情報

機能	Transcription factor for the thyroid-specific expression of the genes exclusively expressed in the thyroid cell type, maintaining the functional differentiation of such cells.
組織特異性	Expressed in the excretory system, thyroid gland and Wilms tumors.
関連疾患	Defects in PAX8 are the cause of congenital hypothyroidism non-goitrous type 2 (CHNG2) [MIM:218700]. CHNG2 is a disease characterized by thyroid dysgenesis, the most frequent cause of congenital hypothyroidism, accounting for 85% of case. The thyroid gland can be completely absent (athyreosis), ectopically located and/or severely hypoplastic. Ectopic thyroid gland is the most frequent malformation, with thyroid tissue being found most often at the base of the tongue.
配列類似性	Contains 1 paired domain.
発生段階	In developing excretory system, during thyroid differentiation and in adult thyroid.
細胞内局在	Nucleus.

画像




PAX8 antibody immunoprecipitates PAX8 protein-DNA in ChIP experiments.

ChIP Sample: HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate;

A. 5 µg preimmune rabbit IgG;

B. 5 µg of ab97477.

The precipitated DNA was detected by PCR with primer set targeting to E2F1 promoter.



Western blot - Anti-PAX8 antibody (ab97477)

All lanes : Anti-PAX8 antibody (ab97477) at 1/10000 dilution

Lane 1 : HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

Lane 2 : DDDDK-tagged PAX8 transfected HEK-293T whole cell lysate


Lysates/proteins at 30 µg per lane.

Secondary

All lanes : HRP-conjugated anti-rabbit IgG antibody

Predicted band size: 48 kDa


10% SDS-PAGE



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PAX8 antibody (ab97477)


Immunohistochemical analysis of paraffin-embedded DLD1 xenograft tissue staining PAX8 protein at nucleolus with ab97477 at 1/500 dilution.

Antigen Retrieval: EDTA based buffer, pH 8.0, 15min.



Immunocytochemistry/ Immunofluorescence - Anti-PAX8 antibody (ab97477)

Immunofluorescence analysis of paraformaldehyde-fixed A549 (human lung carcinoma cell line) whole cells labeling PAX8 with ab97477 at 1/200 dilution. Bottom: Costained with DNA probe stain.



Immunoprecipitation - Anti-PAX8 antibody (ab97477)

ab97477 immunoprecipitates PAX8 protein in IP experiments.

IP Sample: 1000 µg HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate


1- 40 µg HEK-293T whole cell lysate;

2- Control with 2 µg of preimmune rabbit IgG;

3- Immunoprecipitation of PAX8 protein by 2 µg of ab97477.

10% SDS-PAGE


The immunoprecipitated PAX8 protein was detected by ab97477 at 1/1000 dilution, followed by anti-rabbit IgG antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PAX8 antibody (ab97477)

Immunohistochemical analysis of paraffin-embedded mouse liver tissue staining PAX8 protein at nucleus with ab97477 at 1/500 dilution.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.



Western blot - Anti-PAX8 antibody (ab97477)

All lanes : Anti-PAX8 antibody (ab97477) at 1/1000 dilution


Lane 1 : HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

Lane 2 : HEK-293T cell nuclear extract

Lysates/proteins at 30 µg per lane.

Predicted band size: 48 kDa

10% SDS-PAGE



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PAX8 antibody (ab97477)

Immunohistochemical analysis of paraffin-embedded rat thyroid gland tissue staining PAX8 protein at cytosol with ab97477 at 1/500 dilution.

Antigen Retrieval: EDTA based buffer, pH 8.0, 15 min.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors