abcam

Product datasheet

Anti-PBR antibody ab108489

★★★★★ 3 Abreviews 画像数 1

製品の概要

製品名 Anti-PBR antibody

製品の詳細 Rabbit polyclonal to PBR

由来種 Rabbit

 アプリケーション
 適用あり: WB

 種交差性
 交差種: Mouse

交差が予測される動物種: Rat, Horse, Dog, Human, Chimpanzee, Macaque monkey, Gorilla,

Chinese hamster, Orangutan

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール This antibody gave a positive signal in Mouse Brown Adipose tissue.

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquic

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle.

バッファー pH: 7.40

Preservative: 0.02% Sodium azide

Constituent: PBS

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising

agent. If you would like information about the formulation of a specific lot, please contact our

scientific support team who will be happy to help.

精製度 Immunogen affinity purified

ポリ/モノ ポリクローナル

1

アプリケーション


The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab108489の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	**** <u>(2)</u>	Use a concentration of 1 µg/ml. Detects a band of approximately 16 kDa (predicted molecular weight: 19 kDa).

ターゲット情報

機能	Responsible for the manifestation of peripheral-type benzodiazepine recognition sites and is most likely to comprise binding domains for benzodiazepines and isoquinoline carboxamides. May play a role in the transport of porphyrins and heme. Plays a role in the transport of cholesterol across mitochondrial membranes in steroidogenic cells.
組織特異性	Found in many tissue types. Expressed at the highest levels under normal conditions in tissues that synthesize steroids.
配列類似性	Belongs to the TspO/BZRP family.
細胞内局在	Mitochondrion membrane.

画像



Anti-PBR antibody (ab108489) at 1 μ g/ml + P0 Mouse Brown Adipose Tissue Mouse Tissue Lysate at 10 μ g

Secondary

Goat Anti-Rabbit IgG H&L (HRP) preadsorbed (ab97080) at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 19 kDa **Observed band size:** 16 kDa

Additional bands at: 15 kDa. We are unsure as to the identity of

these extra bands.

Exposure time: 3 minutes

The predicted molecular weight of PBR is 19 kDa (SwissProt), however we expect to observe a banding pattern around 16 kDa. Abcam welcomes customer feedback and would appreciate any comments regarding this product and the data presented above.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors