

### Anti-PDGFR alpha antibody [APA5] ab90967

★★★★★ [1 Abreviews](#) [12 References](#) [画像数 1](#)

#### 製品の概要

製品名	Anti-PDGFR alpha antibody [APA5]
製品の詳細	Rat monoclonal [APA5] to PDGFR alpha
由来種	Rat
アプリケーション	<b>適用あり:</b> Flow Cyt
種交差性	<b>交差種:</b> Mouse
免疫原	Fusion protein corresponding to Mouse PDGFR alpha (extracellular). Fusion protein containing the extracellular domain of Mouse PDGFR alpha and the Fc portion of Human IgG1 purified from the culture supernatant of transfected COS1 cells
ポジティブ・コントロール	Flow Cyt: Mouse NIH3T3 cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.20</p> <p>Preservative: 0.09% Sodium azide</p> <p>Constituents: PBS, 0.87% Sodium chloride</p>
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	APA5
アイソタイプ	IgG2a
軽鎖の種類	kappa

## アプリケーション


**The Abpromise guarantee**      **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab90967の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 1 µg for 10 <sup>6</sup> cells. <b>ab18450</b> - Rat monoclonal IgG2a, is suitable for use as an isotype control with this antibody.

## ターゲット情報

機能	Receptor that binds both PDGFA and PDGFB and has a tyrosine-protein kinase activity.
組織特異性	Expressed in primary and metastatic colon tumors and in normal colon tissue. Tumors may express a different isoform to that found in normal tissue.
関連疾患	Note=A chromosomal aberration involving PDGFRA is found in some cases of hypereosinophilic syndrome. Interstitial chromosomal deletion del(4)(q12q12) causes the fusion of FIP1L1 and PDGFRA (FIP1L1-PDGFR).
配列類似性	Belongs to the protein kinase superfamily. Tyr protein kinase family. CSF-1/PDGF receptor subfamily. Contains 5 Ig-like C2-type (immunoglobulin-like) domains. Contains 1 protein kinase domain.
細胞内局在	Membrane.

## 画像



Staining of NIH/3T3 cells with 0.5 µg Rat IgG2a control antibody (open histogram) or 0.5 µg of ab90967 (colored histogram) followed by Biotin Anti-Rat IgG secondary antibody. Total viable cells were analysed.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors