

PE Anti-CCR7 antibody [4B12] ab95669

1 References [画像数 1](#)

製品の概要

製品名	PE Anti-CCR7 antibody [4B12]
製品の詳細	PE Rat monoclonal [4B12] to CCR7
由来種	Rat
標識	PE. Ex: 488nm, Em: 575nm
特異性	ab95669 stains Mouse CCR7-GFP fusion protein-transfected RBL cells
アプリケーション	適用あり: Flow Cyt
種交差性	交差種: Mouse
免疫原	Tissue, cells or virus corresponding to CCR7. CCR7 Rat basophilic leukaemia (RBL)-2H3 transfectants.
ポジティブ・コントロール	C57BL/6, Balb/C and SJL/J splenocytes and thymocytes.
特記事項	Excitation: 488 nm Emission: 575 nm.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 7.20 Preservative: 0.09% Sodium azide Constituents: 0.87% Sodium chloride, PBS
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	4B12
ミエローマ	x63-Ag8.653

アイソタイプ IgG2a
 軽鎖の種類 kappa

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab95669の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 1-5µg for 10 ⁶ cells. ab134674 - Rat monoclonal IgG2a, is suitable for use as an isotype control with this antibody.

ターゲット情報


機能 Receptor for the MIP-3-beta chemokine. Probable mediator of EBV effects on B-lymphocytes or of normal lymphocyte functions.

組織特異性 Expressed in various lymphoid tissues and activated B- and T-lymphocytes, strongly up-regulated in B-cells infected with Epstein-Barr virus and T-cells infected with herpesvirus 6 or 7.

配列類似性 Belongs to the G-protein coupled receptor 1 family.

細胞内局在 Cell membrane.

画像



Staining of C57BL/6 splenocytes with Anti-Mouse CD4 FITC and 0.5 µg of ab95669 with staining carried out at 4 °C (left) or 37 °C (right). Cells in the lymphocyte gate were used for analysis

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors