abcam

Product datasheet

PE Anti-CD22 antibody [MEM-01], prediluted ab27548

画像数1

製品の概要

製品名 PE Anti-CD22 antibody [MEM-01], prediluted

製品の詳細 PE Mouse monoclonal [MEM-01] to CD22, prediluted

由来種 Mouse

標識 PE. Ex: 488nm, Em: 575nm

アプリケーション 適用あり: Flow Cyt 種交差性 交差種: Human

交差が予測される動物種: Non human primates

A

免疫原 Tissue, cells or virus corresponding to CD22. Burkitt's lymphoma cell line Raji

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

バッファー pH: 7.4

Preservative: 0.097% Sodium azide

Constituent: 0.2% BSA

High grade protease free BSA

精製度 IgG fraction ポリ/モノ モノクローナル **クローン名** MEM-01

アイソタイプ lgG1

1


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab27548の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 20µl for 10 ⁶ cells. Use 20µl for 10 ⁶ cells or 100µl of whole blood. ab91357 - Mouse monoclonal lgG1, is suitable for use as an isotype control with this antibody.

ターゲット情報

機能	Mediates B-cell B-cell interactions. May be involved in the localization of B-cells in lymphoid tissues. Binds sialylated glycoproteins; one of which is CD45. Preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site can be masked by cis interactions with sialic acids on the same cell surface. Upon ligand induced tyrosine phosphorylation in the immune response seems to be involved in regulation of B-cell antigen receptor signaling. Plays a role in positive regulation through interaction with Src family tyrosine kinases and may also act as an inhibitory receptor by recruiting cytoplasmic phosphatases via their SH2 domains that block signal transduction through dephosphorylation of signaling molecules.
組織特異性	B-lymphocytes.
配列類似性	Belongs to the immunoglobulin superfamily. SIGLEC (sialic acid binding lg-like lectin) family. Contains 6 lg-like C2-type (immunoglobulin-like) domains. Contains 1 lg-like V-type (immunoglobulin-like) domain.
ドメイン	Contains 4 copies of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.
翻訳後修飾	Phosphorylation of Tyr-762, Tyr-807 and Tyr-822 are involved in binding to SYK, GRB2 and SYK, respectively. Phosphorylation of Tyr-842 is involved in binding to SYK, PLCG2 and PlK3R1/PlK3R2. Phosphorylated on tyrosine residues by LYN.
細胞内局在	Cell membrane.

画像



ab27548 staining CD22 on human peripheral blood cells. Cell population was gated on lymphocytes.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors