

PE Anti-CD45RB antibody [C363.16A] ab25647

1 References [画像数 1](#)

製品の概要

| | |
|--------------|---|
| 製品名 | PE Anti-CD45RB antibody [C363.16A] |
| 製品の詳細 | PE Rat monoclonal [C363.16A] to CD45RB |
| 由来種 | Rat |
| 標識 | PE. Ex: 488nm, Em: 575nm |
| アプリケーション | 適用あり: Flow Cyt |
| 種交差性 | 交差種: Mouse |
| 免疫原 | Tissue, cells or virus corresponding to Mouse CD45RB. Cloned mouse Th cell line |
| エピトープ | ab25647 reacts with an exon B dependent epitope of CD45 glycoprotein. |
| ポジティブ・コントロール | BALB/c mice splenocytes |
| 特記事項 | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

製品の特性

| | |
|--------|--|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at +4°C. Store In the Dark. |
| バッファー | <p>pH: 7.3</p> <p>Preservative: 0.09% Sodium azide</p> <p>Constituents: PBS, 16% Sucrose</p> <p>Also contains a stabilizing agent.</p> |
| 精製度 | Affinity purified |
| ポリ/モノ | モノクローナル |
| クローン名 | C363.16A |
| アイソタイプ | IgG2a |

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab25647の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|--------------------------------------|
| Flow Cyt | | Use 0.2µg for 10 ⁶ cells. |

ターゲット情報

| | |
|-------|--|
| 機能 | Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN. |
| 関連疾患 | Defects in PTPRC are a cause of severe combined immunodeficiency autosomal recessive T-cell-negative/B-cell-positive/NK-cell-positive (T(-)B(+)NK(+) SCID) [MIM:608971]. A form of severe combined immunodeficiency (SCID), a genetically and clinically heterogeneous group of rare congenital disorders characterized by impairment of both humoral and cell-mediated immunity, leukopenia, and low or absent antibody levels. Patients present in infancy recurrent, persistent infections by opportunistic organisms. The common characteristic of all types of SCID is absence of T-cell-mediated cellular immunity due to a defect in T-cell development. Genetic variations in PTPRC are involved in multiple sclerosis susceptibility (MS) [MIM:126200]. MS is a neurodegenerative disorder characterized by the gradual accumulation of focal plaques of demyelination particularly in the periventricular areas of the brain. Peripheral nerves are not affected. Onset usually in third or fourth decade with intermittent progression over an extended period. The cause is still uncertain. |
| 配列類似性 | Belongs to the protein-tyrosine phosphatase family. Receptor class 1/6 subfamily. Contains 2 fibronectin type-III domains. Contains 2 tyrosine-protein phosphatase domains. |
| ドメイン | The first PTPase domain interacts with SKAP1. |
| 翻訳後修飾 | Heavily N- and O-glycosylated. |
| 細胞内局在 | Membrane. Membrane raft. Colocalized with DPP4 in membrane rafts. |

画像



Flow Cytometry - PE Anti-CD45RB antibody
[C363.16A] (ab25647)

BALB/c splenocytes were double-stained with ab25647 (0.1 µg for 10^6 cells) and rat anti-mouse CD8a-FITC. Small lymphocytes were then gated and analyzed on a flow cytometer.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors