

PE Anti-CD80 antibody [16-10A1] ab93507

5 References [画像数 1](#)

製品の概要

製品名	PE Anti-CD80 antibody [16-10A1]
製品の詳細	PE Armenian hamster monoclonal [16-10A1] to CD80
由来種	Armenian hamster
標識	PE. Ex: 488nm, Em: 575nm
アプリケーション	適用あり: Flow Cyt
種交差性	交差種: Mouse
免疫原	Tissue, cells or virus corresponding to Mouse CD80. CHO cell line transfected with Mouse CD80
ポジティブ・コントロール	Activated Mouse splenocyte suspensions
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze. Store undiluted.
バッファー	<p>pH: 7.20</p> <p>Preservative: 0.09% Sodium azide</p> <p>Constituents: 0.87% Sodium chloride, 0.14% Monobasic dihydrogen sodium phosphate, 0.1% Gelatin</p>
精製度	Affinity purified
ポリ/モノ	モノクローナル
クローン名	16-10A1
アイソタイプ	IgG


The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab93507の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use at an assay dependent concentration. ab101026 - Armenian Hamster monoclonal IgG, is suitable for use as an isotype control with this antibody.

ターゲット情報

機能	Involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28 or CTLA-4 to this receptor.
組織特異性	Expressed on activated B-cells, macrophages and dendritic cells.
配列類似性	Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
細胞内局在	Membrane.

画像



Flow Cytometry analysis of C57Bl/6 splenocytes unstimulated (left panel) or stimulated for 3 days with LPS (right panel) labeling CD80 with ab93507 at 0.125µg (solid line). Isotype control - Armenian Hamster IgG (Phycoerythrin) 0.125µg.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors