# abcam

## Product datasheet

## PE Anti-NUR77 antibody [12.14] ab95659

### 画像数1

#### 製品の概要

製品名 PE Anti-NUR77 antibody [12.14]

製品の詳細 PE Mouse monoclonal [12.14] to NUR77

由来種 Mouse

標識 PE. Ex: 488nm, Em: 575nm

アプリケーション 適用あり: Flow Cyt

種交差性 交差種: Mouse

免疫原 Recombinant full length protein corresponding to Mouse NUR77.

ポジティブ・コントロール Mouse thymocytes stimulated with PMA and ionomycin for two hours.

特記事項 The Life Colone industry has been in the order of a new desirible order for a new desirible order.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

#### 製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

**バッファー** pH: 7.20

Preservative: 0.09% Sodium azide

精製度 Protein G purified

ポリ/モノ モノクローナル

**クローン名** 12.14 アイソタイプ IgG1

アプリケーション

## The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab95659の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 1 $\mu$ g for 10 <sup>5-8</sup> cells. in a final volume of 100 $\mu$ L

## ターゲット情報

機能	Orphan nuclear receptor. May act concomitantly with NURR1 in regulating the expression of delayed-early genes during liver regeneration. Binds the NGFIB response element (NBRE) 5'-AAAAGGTCA-3' (By similarity). May inhibit NF-kappa-B transactivation of IL2.	
組織特異性	Fetal muscle and adult liver, brain and thyroid.	
配列類似性	Belongs to the nuclear hormone receptor family. NR4 subfamily.  Contains 1 nuclear receptor DNA-binding domain.	
細胞内局在	Nucleus.	

#### 画像



Flow cytometry images showing staining of unstimulated (left) and 2-hr PMA and lonomycin stimulated (right) C57BL/6 thymocytes with an Anti-Mouse CD4 APC antibody, followed by intracellular staining with 0.5  $\mu$ g of ab95659 using a Foxp3 Staining Buffer Set. Total viable cells were used for analysis.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

## Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors