abcam

Product datasheet

PerCP Anti-CD11a antibody [TP1/40] ab91140

画像数1

製品の概要

製品名 PerCP Anti-CD11a antibody [TP1/40]

製品の詳細 PerCP Mouse monoclonal [TP1/40] to CD11a

由来種 Mouse

標識 PerCP. Ex: 482nm, Em: 675nm

アプリケーション 適用あり: Flow Cyt

種交差性 交差種: Human

免疫原 Tissue, cells or virus corresponding to CD11a. PMA and lonomycin activated T cells blast

(Human)

ポジティブ・コントロール EDTA-anticoagulated blood (lymphocytes, granulocytes, monocytes and macrophages).

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

バッファー pH: 7.2

Preservative: 0.1% Sodium azide

Constituent: 0.164% Sodium phosphate

精製度 Immunogen affinity purified

ポリ/モノ モノクローナル

クローン名 TP1/40

アイソタイプ lgG1

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab91140の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 20µl for 10 ⁶ cells. ab118658 - Mouse monoclonal lgG1, is suitable for use as an isotype control with this antibody.

ターゲット情報

機能 Integrin alpha-L/beta-2 is a receptor for ICAM1, ICAM2, ICAM3 and ICAM4. It is involved in a

 $variety\ of\ immune\ phenomena\ including\ leukocyte-endothelial\ cell\ interaction,\ cytotoxic\ T-cell$

mediated killing, and antibody dependent killing by granulocytes and monocytes.

組織特異性 Leukocytes.

配列類似性 Belongs to the integrin alpha chain family.


Contains 7 FG-GAP repeats.
Contains 1 VWFA domain.

ドメイン The integrin I-domain (insert) is a VWFA domain. Integrins with I-domains do not undergo

protease cleavage.

細胞内局在 Membrane.

画像



Flow cytometry analysis of CD11a expression in normal Human peripheral blood using ab91140 at 20µl/10⁶ cells.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors