abcam

Product datasheet

Anti-Perilipin 3/TIP47 antibody ab118506

画像数1

製品の概要

製品名 Anti-Perilipin 3/TIP47 antibody

製品の詳細 Rabbit polyclonal to Perilipin 3/TIP47

由来種 Rabbit

 アプリケーション
 適用あり: IHC-P

 種交差性
 交差種: Human

交差が予測される動物種: Mouse, Rat

A

免疫原 Synthetic peptide corresponding to Human Perilipin 3/TIP47 (N terminal) conjugated to keyhole

limpet haemocyanin.

ポジティブ・コントロール Human Adrenal tissue.

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term.

バッファー pH: 7.4

Preservative: 0.02% Sodium azide

Constituent: 99% PBS

精製度 Immunogen affinity purified

ポリ/モノ ポリクローナル

アイソタイプ lgG

アプリケーション

1


The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab118506の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

ターゲット情報

機能	Required for the transport of mannose 6-phosphate receptors (MPR) from endosomes to the trans-Golgi network.	
配列類似性	Belongs to the perilipin family.	
細胞内局在	Cytoplasm. Endosome membrane. Lipid droplet. Membrane associated on endosomes. Detected in the envelope and the core of lipid bodies and in lipid sails.	

画像



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Perilipin 3/TIP47 antibody (ab118506)

ab118506, at 5 µg/ml, staining Perilipin 3/TIP47 in formalin-fixed, paraffin-embedded Human Adrenal tissue by Immunohistochemistry using a biotinylated goat anti-rabbit lgG secondary antibody, alkaline phosphatase-streptavidin and chromogen.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors