


### Anti-PGP9.5 antibody [13C4 / I3C4] ab8189

KO 評価済

★★★★☆ 36 Abreviews 115 References 画像数 7

#### 製品の概要

製品名	Anti-PGP9.5 antibody [13C4 / I3C4]
製品の詳細	Mouse monoclonal [13C4 / I3C4] to PGP9.5
由来種	Mouse
アプリケーション	適用あり: ICC, IHC-P, WB
種交差性	交差種: Mouse, Rat, Human 交差が予測される動物種: Sheep, Rabbit, Guinea pig, Dog, Pig, Zebrafish 
免疫原	Full length native protein (purified). This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Wild-type HAP1 whole cell lysate. Human, mouse and rat brain tissue lysate. Rat cortex tissue lysate. SHSY-5Y whole cell lysate. Human spinal cord tissue lysate. IHC-P: Rat pancreas tissue. ICC: Primary rat neurons/glia, DIV14 cells.
特記事項	<p>This antibody labels the neuronal cell bodies and axons in central and peripheral neural system. Small nerve fibers in peripheral tissues, neuroendocrine cells in normal pituitary thyroid, pancreas, and gastrointestinal tract, as well as derived tumors are also stained with this antibody.</p> <p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact <a href="mailto:orders@abcam.com">orders@abcam.com</a>.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40

Preservative: 0.02% Sodium azide  
Constituent: PBS

Some batches contain L-Arginine or BSA as a stabilizing agent. For lot-specific buffer information, please contact our Scientific Support team.

精製度	Protein G purified
一次抗体 備考	This antibody labels the neuronal cell bodies and axons in central and peripheral neural system. Small nerve fibers in peripheral tissues, neuroendocrine cells in normal pituitary thyroid, pancreas, and gastrointestinal tract, as well as derived tumors are also stained with this antibody.
ポリ/モノ	モノクローナル
クローン名	13C4 / I3C4
アイソタイプ	IgG2a


## アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、次のテスト済みアプリケーションにおけるab8189の使用に適用されます**  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ICC		Use a concentration of 5 µg/ml.
IHC-P	★★★★★ (18)	Use a concentration of 0.5 - 5 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
WB	★★★★★ (7)	Use a concentration of 5 µg/ml. Detects a band of approximately 25 kDa (predicted molecular weight: 25 kDa).

## ターゲット情報


機能	Ubiquitin-protein hydrolase involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin. Also binds to free monoubiquitin and may prevent its degradation in lysosomes. The homodimer may have ATP-independent ubiquitin ligase activity.
組織特異性	Found in neuronal cell bodies and processes throughout the neocortex (at protein level). Expressed in neurons and cells of the diffuse neuroendocrine system and their tumors. Weakly expressed in ovary. Down-regulated in brains from Parkinson disease and Alzheimer disease patients.
関連疾患	Parkinson disease 5 Neurodegeneration with optic atrophy, childhood-onset
配列類似性	Belongs to the peptidase C12 family.
翻訳後修飾	O-glycosylated.
細胞内局在	Cytoplasm. Endoplasmic reticulum membrane. About 30% of total UCHL1 is associated with membranes in brain.



Immunocytochemistry - Anti-PGP9.5 antibody [13C4 / I3C4] (ab8189)

ab8189 staining PGP9.5 in primary rat neurons/glia, DIV14 (prepared from E18 rat hippocampal brain area, obtained from Transnetyx Tissue by BrainBits, LLC, cat.no. SDHEP) cells. The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.1% PBS-Tween for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab8189 at 5µg/ml and **ab6046**, Rabbit polyclonal to beta Tubulin - Loading Control. Cells were then incubated with **ab150121**, Goat polyclonal Secondary Antibody to Mouse IgM - mu chain (Alexa Fluor® 488) at 1/1000 dilution (shown in green) and **ab150080**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 594) at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PGP9.5 antibody [13C4 / I3C4] (ab8189)

This image is courtesy of an abreview submitted by Carl Hobbs, King's College London, United Kingdom

IHC-P image of PGP9.5 staining on P5 mouse skin sections using ab8189 (1/1000).

The sections were de-paraffinized and subjected to heat mediated antigen retrieval using citric acid. The sections were then blocked using 1% BSA for 10 mins at 21°C. ab8189 was then incubated for 16 hours at 21°C. The secondary antibody used was Got polyclonal to anti-mouse IgG conjugated to biotin (1/200).




Western blot - Anti-PGP9.5 antibody [13C4 / I3C4] (ab8189)

**Lane 1:** Wild-type HAP1 whole cell lysate (20 µg)

**Lane 2:** PGP9.5 knockout HAP1 whole cell lysate (20 µg)

**Lanes 1 - 4:** Merged signal (red and green). Green - ab8189 observed at 25 kDa. Red - loading control, **ab181602**, observed at 37 kDa.

ab8189 was shown to specifically react with PGP9.5 in wild-type HAP1 cells as signal was lost in PGP9.5 knockout cells. Wild-type and PGP9.5 knockout samples were subjected to SDS-PAGE. ab8189 and **ab181602** (Rabbit anti-GAPDH loading control) were incubated overnight at 4°C at 5 µg/ml and 1/20000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preabsorbed **ab216772** and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preabsorbed **ab216777** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-PGP9.5 antibody [13C4 / I3C4] (ab8189)

**All lanes :** Anti-PGP9.5 antibody [13C4 / I3C4] (ab8189) at 5 µg/ml

**Lane 1 :** Human brain tissue lysate - total protein (**ab29466**)

**Lane 2 :** Brain (Rat) Tissue Lysate

**Lane 3 :** Brain (Mouse) Tissue Lysate

**Lane 4 :** Rat Cortex Tissue Lysate

**Lane 5 :** SHSY-5Y (Human neuroblastoma cell line) Whole Cell Lysate

**Lane 6 :** Human spinal cord tissue lysate - total protein (**ab29188**)

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat Anti-Mouse IgG H&L (HRP) preadsorbed (**ab97040**) at 1/5000 dilution

Developed using the ECL technique.


Performed under reducing conditions.

**Predicted band size:** 25 kDa

**Observed band size:** 25 kDa

**Exposure time:** 1 minute

This blot was produced using a 10% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 5% Bovine Serum Albumin before being incubated with ab8189 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution.




Immunocytochemistry - Anti-PGP9.5 antibody [13C4 / I3C4] (ab8189)

Image courtesy of QBMCellScience

ab8189 (1/20) immunostaining neurons in mouse cortical primary cell culture.






Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PGP9.5 antibody [13C4 / I3C4] (ab8189)

This image is courtesy of an abreview submitted by Carl Hobbs, King's College London, United Kingdom

IHC-P image of PGP9.5 staining on zebrafish brain using ab8189 (1/1000). The sections were subjected to heat mediated antigen retrieval using citric acid. The sections were then blocked using 1% BSA for 10 mins for 21°C. The primary antibody (ab8189) was incubated at a dilution of 1/1000 at 21°C for 16 hours. The secondary antibody used was undiluted goat polyclonal to Mouse IgG conjugated to biotin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PGP9.5 antibody [13C4 / I3C4] (ab8189)

IHC image of PGP9.5 staining in rat pancreas formalin-fixed, paraffin-embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab8189, 0.02µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with hematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors