


Anti-PH3 antibody ab80612

★★★★★ [3 Abreviews](#) [5 References](#) [画像数 2](#)

製品の概要

製品名	Anti-PH3 antibody
製品の詳細	Rabbit polyclonal to PH3
由来種	Rabbit
アプリケーション	適用あり: ICC/IF, WB, IP
種交差性	交差種: Human, Drosophila melanogaster 交差が予測される動物種: Mouse, Rat, Rabbit, Horse, Guinea pig, Dog, Chimpanzee, Ferret, Rhesus monkey, Gorilla, Chinese hamster, Orangutan 
免疫原	Synthetic peptide within Human PH3 aa 700-750. The exact sequence is proprietary. Database link: AAI31773.1
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7 Preservative: 0.09% Sodium azide Constituents: 1.815% Tris, 1.764% Sodium citrate, 0.021% PBS
精製度	Immunogen affinity purified
特記事項(精製)	Purified using an epitope specific to PH3 immobilized on solid support.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab80612の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ICC/IF	★★★★★ (1)	Use at an assay dependent concentration. PubMed: 22019301
WB		1/2000 - 1/10000. Predicted molecular weight: 106 kDa.
IP		Use at 2-5 µg/mg of lysate.

ターゲット情報

機能	Component of the Polycomb group (PcG) multiprotein PRC1 complex, a complex required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility.
配列類似性	Contains 1 FCS-type zinc finger. Contains 1 SAM (sterile alpha motif) domain.
翻訳後修飾	Phosphorylated upon DNA damage, probably by ATM or ATR.
細胞内局在	Nucleus.

画像



All lanes : Anti-PH3 antibody (ab80612) at 0.1 µg/ml

Lane 1 : HeLa whole cell lysate at 50 µg

Lane 2 : HeLa whole cell lysate at 15 µg


Lane 3 : HeLa whole cell lysate at 5 µg

Developed using the ECL technique.

Predicted band size: 106 kDa

Observed band size: 160 kDa

Exposure time: 3 minutes



Immunoprecipitation - Anti-PH3 antibody (ab80612)

Immunoprecipitation of HeLa whole cell lysate using ab80612 at 3µg/mg lysate (lane 1) or a control IgG (lane 2). Bands were detected by Western blot using ab80612 at 1µg/ml. Chemiluminescence with exposure times of 3 minutes.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors