

Anti-Phosphotyrosine antibody [PY20] ab10321

★★★★★ **5 Abreviews** **59 References** 画像数 2

製品の概要

製品名	Anti-Phosphotyrosine antibody [PY20]
製品の詳細	Mouse monoclonal [PY20] to Phosphotyrosine
由来種	Mouse
アプリケーション	適用あり: WB, ICC/IF 適用なし: IHC-P
種交差性	交差種: Species independent
免疫原	Chemical/ Small Molecule corresponding to Phosphotyrosine conjugated to keyhole limpet haemocyanin.
ポジティブ・コントロール	ICC/IF: C2C12 cells treated with 2mM H ₂ O ₂ for 10min. WB: NIH 3T3 treated with Vanadate and PDGF Whole Cell Lysate.
特記事項	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: PBS, 6.97% L-Arginine
精製度	Contains 0.4M Arginine Protein G purified

一次抗体 備考	This is a standard clone used to detect phosphotyrosine.
ポリ/モノ	モノクローナル
クローン名	PY20
アイソタイプ	IgG2b

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab10321の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。


アプリケーション	Abreviews	特記事項
WB	★★★★★ (4)	Use a concentration of 1 µg/ml.
ICC/IF		Use a concentration of 1 µg/ml.

追加情報 Is unsuitable for IHC-P.

ターゲット情報

関連性 The phosphorylation of specific tyrosine residues has been shown to be a primary mechanism of signal transduction during normal mitogenesis, cell cycle progression and oncogenic transformation, its role in other areas such as differentiation and gap junction communication, is a matter of active and ongoing research. Antibodies that specifically recognize phosphorylated tyrosine residues have proved to be invaluable to the study of tyrosine phosphorylated proteins and the biochemical pathways in which they function.

画像




ab10321 staining Phosphotyrosine in C2C12 cells treated with 2mM H₂O₂ for 10mins. The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.1% PBS-Tween for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab10321 at 1µg/ml and **ab6046**, Rabbit polyclonal to beta Tubulin - Loading Control. Cells were then incubated with **ab150117**, Goat polyclonal Secondary Antibody to Mouse IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution (shown in green) and **ab150080**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 594) at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Also suitable in cells fixed with 100% methanol (5 min).

Image was acquired with a confocal microscope (Leica-

Microsystems TCS SP8) and a single confocal section is shown.



All lanes : Anti-Phosphotyrosine antibody [PY20] (ab10321) at 1 μ g/ml

Lane 1 : NIH 3T3 (Mouse embryonic fibroblast cell line) Whole Cell Lysate

Lane 2 : NIH 3T3 treated with Vanadate and PDGF Whole Cell Lysate

Lysates/proteins at 5 μ g per lane.

Secondary

All lanes : Goat polyclonal to Mouse IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Exposure time: 1 minute

Cells were serum starved overnight and then incubated at room temperature for 10mins in a final concentration of 1mM sodium vanadate. PDGF was then added at a final concentration of 5ng/ml and cells were incubated at 37°C for 30mins. Vanadate inhibits endogenous phosphatases and PDGF stimulates phosphorylation. Western blots of NIH 3T3 cell lysates treated with vanadate and PDGF show an array of phosphorylated tyrosine compared to controls.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors