


Anti-PIAS1 antibody [EPR2580(2)] ab109388

リコンビナント **RabMAb**


★★★★★ **3 Abreviews** **16 References** 画像数 6

製品の概要

製品名	Anti-PIAS1 antibody [EPR2580(2)]
製品の詳細	Rabbit monoclonal [EPR2580(2)] to PIAS1
由来種	Rabbit
アプリケーション	適用あり: ICC/IF, WB, IHC-P 適用なし: IP
種交差性	交差種: Human 交差が予測される動物種: Mouse, Rat 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	Human colon tissue. HepG2, Jurkat, 293T, Daudi cell lysates. ICC/IF: HeLa cells
特記事項	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
解離定数 (K _D 値)	K _D = 2.10 x 10 ⁻¹¹ M



バッファー	pH: 7.20 Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR2580(2)
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab109388の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF	★★★★★ (1)	1/500.
WB	★★★★★ (2)	1/1000 - 1/10000. Predicted molecular weight: 72 kDa.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol. Heat up to 98 degrees C, below boiling, and then let cool for 10-20 min.

追加情報 Is unsuitable for IP.

ターゲット情報


機能	Functions as an E3-type small ubiquitin-like modifier (SUMO) ligase, stabilizing the interaction between UBE2I and the substrate, and as a SUMO-tethering factor. Plays a crucial role as a transcriptional coregulation in various cellular pathways, including the STAT pathway, the p53 pathway and the steroid hormone signaling pathway. In vitro, binds A/T-rich DNA. The effects of this transcriptional coregulation, transactivation or silencing, may vary depending upon the biological context. Together with PRMT1, may repress STAT1 transcriptional activity, in the late phase of interferon gamma (IFN-gamma) signaling.
組織特異性	Expressed in numerous tissues with highest level in testis.
パスウェイ	Protein modification; protein sumoylation.
配列類似性	Belongs to the PIAS family. Contains 1 PINIT domain. Contains 1 SAP domain. Contains 1 SP-RING-type zinc finger.
ドメイン	The LXXLL motif is a transcriptional coregulator signature. The SP-RING-type domain is required for promoting EKLF sumoylation.
翻訳後修飾	Sumoylated. Dimethylated by PRMT1 at Arg-303 in the late phase of interferon gamma (IFN-gamma) signaling, leading to preferential interaction with STAT1 and thus resulting in release of STAT1

from its target gene.

細胞内局在

Nucleus speckle. Interaction with CSRP2 may induce a partial redistribution along the cytoskeleton.

画像



Lane 1 : Anti-PIAS1 antibody [EPR2580(2)] (ab109388) at 1/500 dilution

Lane 2 : Anti-GST antibody [EPR4236] ([ab111947](#)) at 1/2000 dilution

All lanes : Recombinant Human PIAS1 protein ([ab152888](#))

Lysates/proteins at 0.01 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 72 kDa


Observed band size: 98 kDa

Exposure time:

Lane 1: 10 seconds.


Lane 2: 1 minute.

Blocking and dilution buffer: 5% NFDM /TBST.



Immunocytochemistry/ Immunofluorescence - Anti-PIAS1 antibody [EPR2580(2)] (ab109388)

Immunocytochemistry/ Immunofluorescence analysis of HeLa (human cervix adenocarcinoma epithelial cell) cells labeling PIAS1 with purified ab109388 at 1/500 dilution (4.6 µg/mL). Cells were fixed in 100% Methanol. Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1/1000 (2 µg/mL) dilution. DAPI (blue) was used as nuclear counterstain.



Western blot - Anti-PIAS1 antibody [EPR2580(2)] (ab109388)

All lanes : Anti-PIAS1 antibody [EPR2580(2)] (ab109388) at 1/1000 dilution

Lane 1 : HepG2 cell lysates


Lane 2 : Jurkat cell lysates

Lane 3 : HEK-293 cell lysates

Lane 4 : Daudi cell lysates

Lysates/proteins at 10 µg per lane.


Predicted band size: 72 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PIAS1 antibody [EPR2580(2)] (ab109388)

Immunohistochemical analysis of paraffin-embedded Human colon tissue using 1/100 ab109388.

Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.



SPR Scanning - Anti-PIAS1 antibody [EPR2580(2)]
(ab109388)

Equilibrium dissociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-PIAS1 antibody [EPR2580(2)] (ab109388)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors