

Anti-PMS2 antibody [EPR3947] ab110638

KO 評価済 リコンビナント RabMAb


★★★★☆ 3 Abreviews 30 References 画像数 12

製品の概要

| | |
|--------------|--|
| 製品名 | Anti-PMS2 antibody [EPR3947] |
| 製品の詳細 | Rabbit monoclonal [EPR3947] to PMS2 |
| 由来種 | Rabbit |
| アプリケーション | 適用あり: Flow Cyt (Intra), WB, IP, IHC-P, ICC/IF |
| 種交差性 | 交差種: Human |
| 免疫原 | Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. |
| ポジティブ・コントロール | WB: Hap1 and HeLa cell lysates. IP: HeLa whole cell lysate. |
| 特記事項 | <p>To see more of the key markers and tools you need to study the hallmarks of cancer, including genome instability and mutation, please visit the following page.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> |

製品の特性

| | |
|-------------------------|--|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C. |
| 解離定数 (K _D 値) | K _D = 1.50 x 10 ⁻¹⁰ M |



| | |
|-------|----------|
| バッファー | pH: 7.20 |
|-------|----------|

| | |
|--------|--|
| | Preservative: 0.01% Sodium azide |
| | Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA |
| 精製度 | Protein A purified |
| ポリ/モノ | モノクローナル |
| クローン名 | EPR3947 |
| アイソタイプ | IgG |

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab110638の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項 |
|------------------|-----------|--|
| Flow Cyt (Intra) | | 1/160. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. |
| WB | ★★★★★ (2) | 1/10000. Detects a band of approximately 110 kDa (predicted molecular weight: 96 kDa). |
| IP | | 1/10 - 1/100. |
| IHC-P | | 1/600. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. It is recommended to trial a dilution factor from 1:50 with IHC-P as the staining system sensitivity can be variable |
| ICC/IF | ★★★★★ (1) | 1/200. |

ターゲット情報

| | |
|------|---|
| 機能 | Component of the post-replicative DNA mismatch repair system (MMR). Heterodimerizes with MLH1 to form MutL alpha. DNA repair is initiated by MutS alpha (MSH2-MSH6) or MutS beta (MSH2-MSH6) binding to a dsDNA mismatch, then MutL alpha is recruited to the heteroduplex. Assembly of the MutL-MutS-heteroduplex ternary complex in presence of RFC and PCNA is sufficient to activate endonuclease activity of PMS2. It introduces single-strand breaks near the mismatch and thus generates new entry points for the exonuclease EXO1 to degrade the strand containing the mismatch. DNA methylation would prevent cleavage and therefore assure that only the newly mutated DNA strand is going to be corrected. MutL alpha (MLH1-PMS2) interacts physically with the clamp loader subunits of DNA polymerase III, suggesting that it may play a role to recruit the DNA polymerase III to the site of the MMR. Also implicated in DNA damage signaling, a process which induces cell cycle arrest and can lead to apoptosis in case of major DNA damages. |
| 関連疾患 | Defects in PMS2 are the cause of hereditary non-polyposis colorectal cancer type 4 (HNPCC4) [MIM:600259]. Mutations in more than one gene locus can be involved alone or in combination in the production of the HNPCC phenotype (also called Lynch syndrome). Most families with clinically recognized HNPCC have mutations in either MLH1 or MSH2 genes. HNPCC is an |

autosomal, dominantly inherited disease associated with marked increase in cancer susceptibility. It is characterized by a familial predisposition to early onset colorectal carcinoma (CRC) and extra-colonic cancers of the gastrointestinal, urological and female reproductive tracts. HNPCC is reported to be the most common form of inherited colorectal cancer in the Western world, and accounts for 15% of all colon cancers. Cancers in HNPCC originate within benign neoplastic polyps termed adenomas. Clinically, HNPCC is often divided into two subgroups. Type I: hereditary predisposition to colorectal cancer, a young age of onset, and carcinoma observed in the proximal colon. Type II: patients have an increased risk for cancers in certain tissues such as the uterus, ovary, breast, stomach, small intestine, skin, and larynx in addition to the colon. Diagnosis of classical HNPCC is based on the Amsterdam criteria: 3 or more relatives affected by colorectal cancer, one a first degree relative of the other two; 2 or more generation affected; 1 or more colorectal cancers presenting before 50 years of age; exclusion of hereditary polyposis syndromes. The term 'suspected HNPCC' or 'incomplete HNPCC' can be used to describe families who do not or only partially fulfill the Amsterdam criteria, but in whom a genetic basis for colon cancer is strongly suspected. Defects in PMS2 are a cause of mismatch repair cancer syndrome (MMRCS) [MIM:276300]; also known as Turcot syndrome or brain tumor-polyposis syndrome 1 (BTPS1). MMRCS is an autosomal dominant disorder characterized by malignant tumors of the brain associated with multiple colorectal adenomas. Skin features include sebaceous cysts, hyperpigmented and cafe au lait spots.


配列類似性

Belongs to the DNA mismatch repair mutL/hexB family.

細胞内局在

Nucleus.

画像



All lanes : Anti-PMS2 antibody [EPR3947] (ab110638) at 1/10000 dilution (Purified)


Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

Lane 2 : Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 96 kDa




Flow Cytometry (Intracellular) - Anti-PMS2 antibody
[EPR3947] (ab110638)

Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labelling PMS2 with Purified ab110638 at 1:160 dilution (10 µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150081](#)) secondary antibody was used at 1:2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabelled control - Cell without incubation with primary antibody and secondary antibody (Blue).




Immunocytochemistry/ Immunofluorescence - Anti-PMS2 antibody [EPR3947] (ab110638)

Immunocytochemistry analysis of SH-SY5Y (Human neuroblastoma epithelial cell) cells labeling PMS2 with Purified ab110638 at 1:200 dilution (8 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PMS2 antibody [EPR3947] (ab110638)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast tissue sections labeling PMS2 with Purified ab110638 at 1:600 (2.71 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.



Western blot - Anti-PMS2 antibody [EPR3947] (ab110638)

All lanes : Anti-PMS2 antibody [EPR3947] (ab110638) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : PMS2 knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


Predicted band size: 96 kDa

Observed band size: 120 kDa

Lanes 1- 2: Merged signal (red and green). Green - ab110638 observed at 120 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) observed at 37 kDa.

ab110638 was shown to react with PMS2 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line

ab261776 (knockout cell lysate **ab257142**) was used. Wild-type HeLa and PMS2 knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab110638 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-PMS2 antibody [EPR3947]
(ab110638)

All lanes : Anti-PMS2 antibody [EPR3947] (ab110638) at 1/1000 dilution

Lane 1 : Wild-type HAP1 whole cell lysate


Lane 2 : PMS2 knockout HAP1 whole cell lysate

Lysates/proteins at 30 µg per lane.

Predicted band size: 96 kDa

Lanes 1-2: Merged signal (red and green). Green - ab110638 observed at 120 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

ab110638 was shown to specifically react with PMS2 in wild-type HAP1 cells. No band was observed when PMS2 knockout samples were used. Wild-type and PMS2 knockout samples were subjected to SDS-PAGE. Ab110638 and **ab8245** (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PMS2 antibody [EPR3947] (ab110638)

Immunohistochemical analysis of paraffin-embedded Human colon tissue labelling PMS2 with ab110638 at 1/100 followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**). Positive staining on human colon. The section was incubated with ab110638 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PMS2 antibody [EPR3947] (ab110638)

Immunohistochemical analysis of paraffin-embedded Human colon cancer tissue labelling PMS2 with ab110638 at 1/100 followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**). Positive staining on human colon cancer. The section was incubated with ab110638 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunoprecipitation - Anti-PMS2 antibody
[EPR3947] (ab110638)

Purified ab110638 at 1/50 dilution (2µg) immunoprecipitating PMS2 in HeLa whole cell lysate.

Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10µg

Lane 2 (+): ab110638 + HeLa whole cell lysate.

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab110638 in HeLa whole cell lysate.


VeriBlot for IP Detection Reagent (HRP) (**ab131366**) (1/5000 dilution) was used for Western blotting.

Blocking Buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM/TBST.

Observed band size: 110 kDa

Lower bands are degradation bands and fresh lysate is recommended.



Immunoprecipitation - Anti-PMS2 antibody
[EPR3947] (ab110638)

Purified ab110638 at 1/50 dilution (2µg) immunoprecipitating PMS2 in HeLa whole cell lysate.

Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10µg

Lane 2 (+): ab110638 + HeLa whole cell lysate.

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab110638 in HeLa whole cell lysate.

VeriBlot for IP Detection Reagent (HRP) (**ab131366**) (1/5000 dilution) was used for Western blotting.

Blocking Buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM/TBST.

Observed band size: 110 kDa

Lysate were made freshly and used in IP test immediately to minimize protein degradation. Incubation time was 2h.

| Tissue Microarray (TMA) data for ab110638 | | | |
|---|---|--------------------------------------|---|
| Normal tissue samples | | Malignant tissue samples | |
| Human cardiac muscle | x | Human placenta | x |
| Human cerebrum | ✓ | Human skeletal muscle | x |
| Human colon | ✓ | Human skin | ✓ |
| Human endometrium | ✓ | Human spleen | ✓ |
| Human kidney | ✓ | Human stomach | ✓ |
| Human liver | x | Human testis | ✓ |
| Human lung | x | Human thyroid | ✓ |
| Human mammary gland | ✓ | Human tonsil | ✓ |
| Human pancreas | ✓ | | |
| | | Clear cell carcinoma of human kidney | x |
| | | Human bladder cancer | ✓ |
| | | Human breast carcinoma | ✓ |
| | | Human cervical carcinoma | ✓ |
| | | Human colon carcinoma | ✓ |
| | | Human endometrial carcinoma | ✓ |
| | | Human gastric adenocarcinoma | ✓ |
| | | Human glioma | ✓ |
| | | Human hepatocellular carcinoma | ✓ |
| | | Human lung carcinoma | ✓ |
| | | Human melanoma | x |
| | | Human ovarian carcinoma | ✓ |
| | | Human prostatic hyperplasia | x |
| | | Human thyroid carcinoma | ✓ |

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PMS2 antibody [EPR3947] (ab110638)

Tissue Microarrays stained for "Anti-PMS2 antibody [EPR3947]" using "ab110638" in immunohistochemical analysis. This table provides a detailed overview of positive (tick mark) and negative (cross mark) staining per sample type tested. The sections were pre-treated using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) for 20 minutes. The sections were incubated with ab110638 for 30 mins at room temperature followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**). The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Research with confidence
Consistent and reproducible results

Long-term and scalable supply
Recombinant technology

Success from the first experiment
Confirmed specificity

Ethical standards compliant
Animal-free production

Anti-PMS2 antibody [EPR3947] (ab110638)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors