

Anti-Polyethylene glycol antibody [26A04] - BSA and Azide free ab94764

★★★★★ [1 Abreviews](#) [7 References](#) [画像数 2](#)

製品の概要

製品名	Anti-Polyethylene glycol antibody [26A04] - BSA and Azide free
製品の詳細	Rat monoclonal [26A04] to Polyethylene glycol - BSA and Azide free
由来種	Rat
特異性	ab94764 recognizes linear 20 kDa PEG, linear 5 kDa PEG, linear 600 Da PEG and 10 kDa branched PEG. It also binds to BSA conjugated PEG by western blot.
アプリケーション	適用あり: WB, Indirect ELISA
種交差性	交差種: Species independent
免疫原	5 kDa PEG conjugated to KLH by a MBS linker.
ポジティブ・コントロール	PEGylated protein
特記事項	<p>Tween 20 usage: Tween 20 is a detergent that can interfere with ELISA and potentially other applications (often used in antibody dilution buffers, such as TBST) when using anti-PEG products. This is also the same case for other polyoxyethylene detergents. Therefore, it is recommended to avoid Tween 20 and other polyoxyethylene detergents when performing ELISA using anti-PEG products.</p> <p><u>Learn more about PEG RabMAb primary antibodies and kit - PEG RabMAb Product Portal Page</u></p>

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.

バッファー	Constituent: Tris buffered saline
キャリア・フリー	はい
特記事項 (精製)	Ceramic hydroxyapatite (CHT) chromatography from in vitro production.
ポリ/モノ	モノクローナル
クローン名	26A04
アイソタイプ	IgM


アプリケーション

The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab94764の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Predicted molecular weight: 1 kDa.
Indirect ELISA		Use a concentration of 2 µg/ml.

ターゲット情報

画像



Indirect ELISA analysis of a PEG panel using ab94764.



Anti-Polyethylene glycol antibody [26A04] - BSA and Azide free (ab94764) + PEGylated protein

Predicted band size: 1 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors