

Anti-PPAR gamma antibody ab59256

★★★★★ [2 Abreviews](#) [60 References](#) [画像数 3](#)

製品の概要

製品名	Anti-PPAR gamma antibody
製品の詳細	Rabbit polyclonal to PPAR gamma
由来種	Rabbit
アプリケーション	適用あり: WB, ICC/IF, IHC-P
種交差性	交差種: Human
免疫原	Synthetic non-phosphopeptide derived from human PPAR gamma around the phosphorylation site of serine 112 (P-A-S ^P -P-P).
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	<p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituents: PBS, 50% Glycerol, 0.87% Sodium chloride</p>
精製度	Without Mg+2 and Ca+2
特記事項 (精製)	Immunogen affinity purified
ポリ/モノ	ab59256 was affinity purified from rabbit antiserum by affinity chromatography using epitope specific immunogen.
アイソタイプ	ポリクローナル
	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab59256の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		1/500 - 1/1000. Predicted molecular weight: 57 kDa.
ICC/IF		Use a concentration of 5 µg/ml.
IHC-P	★★★★★ (2)	1/50 - 1/100.

ターゲット情報

機能	Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the receptor binds to a promoter element in the gene for acyl-CoA oxidase and activates its transcription. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids. Key regulator of adipocyte differentiation and glucose homeostasis.
組織特異性	Highest expression in adipose tissue. Lower in skeletal muscle, spleen, heart and liver. Also detectable in placenta, lung and ovary.
関連疾患	<p>Note=Defects in PPARG can lead to type 2 insulin-resistant diabetes and hypertension. PPARG mutations may be associated with colon cancer.</p> <p>Defects in PPARG may be associated with susceptibility to obesity (OBESITY) [MIM:601665]. It is a condition characterized by an increase of body weight beyond the limitation of skeletal and physical requirements, as the result of excessive accumulation of body fat.</p> <p>Defects in PPARG are the cause of familial partial lipodystrophy type 3 (FPLD3) [MIM:604367]. Familial partial lipodystrophies (FPLD) are a heterogeneous group of genetic disorders characterized by marked loss of subcutaneous (sc) fat from the extremities. Affected individuals show an increased preponderance of insulin resistance, diabetes mellitus and dyslipidemia.</p> <p>Genetic variations in PPARG can be associated with susceptibility to glioma type 1 (GLM1) [MIM:137800]. Gliomas are central nervous system neoplasms derived from glial cells and comprise astrocytomas, glioblastoma multiforme, oligodendrogliomas, and ependymomas.</p> <p>Note=Polymorphic PPARG alleles have been found to be significantly over-represented among a cohort of American patients with sporadic glioblastoma multiforme suggesting a possible contribution to disease susceptibility.</p>
配列類似性	<p>Belongs to the nuclear hormone receptor family. NR1 subfamily.</p> <p>Contains 1 nuclear receptor DNA-binding domain.</p>
細胞内局在	Nucleus.

画像




All lanes : Anti-PPAR gamma antibody (ab59256) at 1/500 dilution


Lane 1 : HUVEC (human umbilical vein endothelial cell line) whole cell lysate

Lane 2 : HUVEC (blocked with immunizing peptide), whole cell lysate

Predicted band size: 57 kDa



ab59256, at 1/50 dilution, staining PPAR gamma in paraffin embedded human placenta tissue by Immunohistochemistry in the absence or presence of the immunising peptide.



Immunocytochemistry/ Immunofluorescence - Anti-PPAR gamma antibody (ab59256)

ICC/IF image of ab59256 stained MCF7 cells. The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab59256, 5µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors