

### Anti-pro Caspase-7 antibody [Y33] ab32067

KO 評価済 リコンビナント RabMAb

2 References 画像数 4

#### 製品の概要

製品名	Anti-pro Caspase-7 antibody [Y33]
製品の詳細	Rabbit monoclonal [Y33] to pro Caspase-7
由来種	Rabbit
特異性	The antibody only recognizes the pro-form of Caspase-7. It does not react with the cleaved forms (active enzyme) of Caspase-7.
アプリケーション	<b>適用あり:</b> WB, IHC-P <b>適用なし:</b> Flow Cyt or IP
種交差性	<b>交差種:</b> Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	Jurkat cell lysate and human lung squamous carcinoma.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 49% PBS, 50% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified

ポリ/モノ	モノクローナル
クローン名	Y33
アイソタイプ	IgG

## アプリケーション

**The Abpromise guarantee** **Abpromise保証は、次のテスト済みアプリケーションにおけるab32067の使用に適用されます**  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB		1/1000. Detects a band of approximately 34 kDa (predicted molecular weight: 34 kDa).
IHC-P		1/50. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

**追加情報** Is unsuitable for Flow Cyt or IP.

## ターゲット情報

<b>機能</b>	Involved in the activation cascade of caspases responsible for apoptosis execution. Cleaves and activates sterol regulatory element binding proteins (SREBPs). Proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at a '216-Asp-Gly-217' bond. Overexpression promotes programmed cell death.
<b>組織特異性</b>	Highly expressed in lung, skeletal muscle, liver, kidney, spleen and heart, and moderately in testis. No expression in the brain.
<b>配列類似性</b>	Belongs to the peptidase C14A family.
<b>翻訳後修飾</b>	Cleavages by granzyme B or caspase-10 generate the two active subunits. Propeptide domains can also be cleaved efficiently by caspase-3. Active heterodimers between the small subunit of caspase-7 and the large subunit of caspase-3, and vice versa, also occur.
<b>細胞内局在</b>	Cytoplasm.

## 画像



Western blot - Anti-pro Caspase-7 antibody [Y33]  
(ab32067)

**Lane 1:** Wild type HAP1 whole cell lysate (20 µg)

**Lane 2:** Staurosporin treated HAP1 whole cell lysate (20 µg)

**Lane 3:** CASP7 knockout HAP1 whole cell lysate (20 µg)


**Lane 4:** Staurosporin treated CASP7 knockout HAP1 whole cell lysate (20 µg)

**Lane 5:** HeLa whole cell lysate (20 µg)

**Lane 6:** Staurosporin treated HeLa whole cell lysate (20 µg)

**Lanes 1 - 6:** Merged signal (red and green). Green - ab32067 observed at 38 kDa. Red - loading control, **ab18058**, observed at 130 kDa.


ab32067 was shown to recognize pro Caspase 7 when HAP1 + pro Caspase 7 knockout samples were used, along with additional cross-reactive bands. Wild-type and the pro Caspase 7 knockout samples were subjected to SDS-PAGE. Ab32067 and **ab18058** (Mouse anti-Vinculin loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/10000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-pro Caspase-7 antibody [Y33] (ab32067)

Immunohistochemical analysis of paraffin-embedded human lung squamous carcinoma using ab32067 at 1/50 dilution.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-pro Caspase-7 antibody [Y33] (ab32067)

Anti-pro Caspase-7 antibody [Y33] (ab32067) at 1/1000 dilution + Jurkat cell lysate

**Predicted band size:** 34 kDa

**Observed band size:** 34 kDa

#### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-pro Caspase-7 antibody [Y33] (ab32067)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors