

Anti-RAGE antibody [1C1] ab54741

★★★★★ [5 Abreviews](#) [19 References](#) [画像数 5](#)

製品の概要

製品名	Anti-RAGE antibody [1C1]
製品の詳細	Mouse monoclonal [1C1] to RAGE
由来種	Mouse
アプリケーション	適用あり: WB, IP, IHC-P
種交差性	交差種: Human, Recombinant fragment
免疫原	Recombinant full length protein (GST-tag) corresponding to Human RAGE aa 23-405. Full length mature chain, without signal peptide. Database link: Q15109
ポジティブ・コントロール	WB: Human kidney tissue lysate. RAGE transfected HEK-293T cell lysate. Recombinant RAGE protein. Flow Cytometry: Jurkat cells. IHC-P: Human lung tissue.
特記事項	<p>This product was changed from ascites to tissue culture supernatant on 13th Feb 2019. Please note that the dilutions may need to be adjusted accordingly. If you have any questions, please do not hesitate to contact our scientific support team.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.40 Constituent: PBS
精製度	In 1x PBS Tissue culture supernatant

ポリ/モノ	モノクローナル
クローン名	1C1
アイソタイプ	IgG2a
軽鎖の種類	kappa

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab54741の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (3)	Use at an assay dependent concentration. Predicted molecular weight: 43 kDa. This antibody has only been tested in WB against the recombinant fragment used as immunogen. We have no data on the detection of endogenous protein in Western blot
IP		Use at an assay dependent concentration.
IHC-P	★★★★★ (1)	Use at an assay dependent concentration. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

ターゲット情報

機能	Mediates interactions of advanced glycosylation end products (AGE). These are nonenzymatically glycosylated proteins which accumulate in vascular tissue in aging and at an accelerated rate in diabetes. Acts as a mediator of both acute and chronic vascular inflammation in conditions such as atherosclerosis and in particular as a complication of diabetes. AGE/RAGE signaling plays an important role in regulating the production/expression of TNF-alpha, oxidative stress, and endothelial dysfunction in type 2 diabetes. Interaction with S100A12 on endothelium, mononuclear phagocytes, and lymphocytes triggers cellular activation, with generation of key proinflammatory mediators. Interaction with S100B after myocardial infarction may play a role in myocyte apoptosis by activating ERK1/2 and p53/TP53 signaling (By similarity). Receptor for amyloid beta peptide. Contributes to the translocation of amyloid-beta peptide (ABPP) across the cell membrane from the extracellular to the intracellular space in cortical neurons. ABPP-initiated RAGE signaling, especially stimulation of p38 mitogen-activated protein kinase (MAPK), has the capacity to drive a transport system delivering ABPP as a complex with RAGE to the intraneuronal space.
組織特異性	Endothelial cells.
配列類似性	Contains 2 Ig-like C2-type (immunoglobulin-like) domains. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
細胞内局在	Secreted and Cell membrane.

画像




Western blot - Anti-RAGE antibody [1C1] (ab54741)

Anti-RAGE antibody [1C1] (ab54741) at 1 µg/ml + Human kidney tissue lysate

Developed using the ECL technique.

Predicted band size: 43 kDa

This image was generated using the ascites version of the product.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RAGE antibody [1C1] (ab54741)

ab54741 (1 µg/ml) staining RAGE in human lung using an automated system (DAKO Autostainer Plus). Using this protocol there is membrane staining throughout the alveoli.

Sections were rehydrated and antigen retrieved with the Dako 3 in 1 AR buffer citrate pH 6.0 in a DAKO PT link. Slides were peroxidase blocked in 3% H₂O₂ in methanol for 10 minutes. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 minutes and detected with Dako envision flex amplification kit for 30 minutes. Colorimetric detection was completed with Diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that, for manual staining, optimization of primary antibody concentration and incubation time is recommended. Signal amplification may be required.

This image was generated using the ascites version of the product.



Western blot - Anti-RAGE antibody [1C1] (ab54741)

All lanes : Anti-RAGE antibody [1C1] (ab54741) at 1 µg/ml


Lane 1 : HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) RAGE transfected cell lysate

Lane 2 : HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) non-transfected cell lysate

Developed using the ECL technique.

Predicted band size: 43 kDa

This image was generated using the ascites version of the product.



Western blot - Anti-RAGE antibody [1C1] (ab54741)

Anti-RAGE antibody [1C1] (ab54741) at 1 µg/ml + tagged recombinant protein immunogen

Developed using the ECL technique.

Predicted band size: 43 kDa

Predicted band size of immunogen is 68 kDa

This image was generated using the ascites version of the product.



Immunoprecipitation - Anti-RAGE antibody [1C1] (ab54741)

Immunoprecipitation of RAGE transfected lysate using ab54741 and Protein A Magnetic Bead. Immunoblotted with a rabbit polyclonal anti-RAGE antibody.

This image was generated using the ascites version of the product.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors