



Anti-RHCG antibody ab106837

画像数 1

製品の概要

製品名	Anti-RHCG antibody
製品の詳細	Goat polyclonal to RHCG
由来種	Goat
アプリケーション	適用あり: WB
種交差性	交差種: Human
免疫原	Synthetic peptide: ERQNSVYQSD by a Cysteine residue linker, corresponding to internal sequence amino acids 209-218 of Human RHCG (NP_057405.1).
	 Run BLAST with  Run BLAST with
ポジティブ・コントロール	Human testis tissue lysate.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: Tris buffered saline, 0.5% BSA
精製度	Immunogen affinity purified
特記事項 (精製)	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab106837の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 - 3 µg/ml. Predicted molecular weight: 53 kDa. 1 hour primary incubation is recommended for this product.

ターゲット情報

機能	Functions as an electroneutral and bidirectional ammonium transporter. May regulate transepithelial ammonia secretion.
組織特異性	Expressed in brain, testis, placenta, pancreas, esophagus and prostate. Expressed in squamous epithelial tissues (at protein level). According to PubMed:11062476, specifically expressed in kidney.
配列類似性	Belongs to the ammonium transporter (TC 2.A.49) family. Rh subfamily.
発生段階	Specifically expressed in fetal kidney.
翻訳後修飾	N-glycosylated.
細胞内局在	Apical cell membrane. Also detected at the basolateral membrane and in subapical vesicles.

画像



Anti-RHCG antibody (ab106837) at 1 µg/ml + Human Testis lysate
in RIPA buffer. at 35 µg

Predicted band size: 53 kDa

Primary incubation was 1 hour. Detected by chemiluminescence

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors