

Anti-RNF185 antibody ab73118

画像数 1

製品の概要

製品名	Anti-RNF185 antibody
製品の詳細	Mouse polyclonal to RNF185
由来種	Mouse
アプリケーション	適用あり: WB
種交差性	交差種: Human
免疫原	Recombinant full length protein within Human RNF185. The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please contact our Scientific Support team to discuss your requirements.
ポジティブ・コントロール	RNF185 transfected 293T cell lysate
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.4 Constituents: 0.1136% Dibasic monohydrogen sodium phosphate, Whole serum, 0.812% Sodium chloride, 0.0225% Potassium chloride, 0.0204% Monobasic dihydrogen potassium phosphate
精製度	Purified IgA
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab73118の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		1/500 - 1/1000. Predicted molecular weight: 20 kDa.

ターゲット情報

機能	E3 ubiquitin-protein ligase that regulates selective mitochondrial autophagy by mediating 'Lys-63'-linked polyubiquitination of BNIP1. Responsible for the cotranslational ubiquitination and degradation of CFTR in the ERAD pathway. Preferentially associates with the E2 enzymes UBE2J1 and UBE2J2.
パスウェイ	Protein modification; protein ubiquitination.
配列類似性	Contains 1 RING-type zinc finger.
ドメイン	The RING-type zinc finger domain is responsible for E3 ubiquitin ligase activity.
細胞内局在	Mitochondrion outer membrane. Endoplasmic reticulum membrane.

画像



All lanes : Anti-RNF185 antibody (ab73118) at 1/500 dilution

Lane 1 : RNF185 transfected 293T cell lysate

Lane 2 : non transfected lysate

Lysates/proteins at 25 µg per lane.

Secondary

All lanes : goat anti-mouse IgG (H&L)-HRP conjugate at 1/2500 dilution

Predicted band size: 20 kDa

Observed band size: 21 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors