

Anti-S tag antibody ab183674

4 References [画像数 2](#)

製品の概要

製品名	Anti-S tag antibody
製品の詳細	Rabbit polyclonal to S tag
由来種	Rabbit
アプリケーション	適用あり: WB, IP
種交差性	交差種: Species independent
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	S tag transfected 293T cell lysate.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.00</p> <p>Preservative: 0.025% Proclin 300</p> <p>Constituents: 79% PBS, 20% Glycerol (glycerin, glycerine)</p>
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab183674の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		1/1000 - 1/10000.
IP		Use at an assay dependent concentration. Use 2.5µg.

ターゲット情報

関連性 S tag is an epitope tag composed of a 15 residue peptide, KETAAAKFERQHMS, derived from the pancreatic ribonuclease A.

画像




All lanes : Anti-S tag antibody (ab183674) at 1/5000 dilution

Lane 1 : 293T whole cell lysate

Lane 2 : S tag transfected 293T whole cell lysate

Lysates/proteins at 30 µg per lane.

12% SDS-PAGE



S tag antibody ab183674 immunoprecipitates S tag protein in IP experiments. IP Sample: 650 µg whole cell lysate of S tag protein expressing HeLa cells (1), 20 µg whole cell lysate of transfected HeLa cells (2), Control with 2.5 µg of preimmune rabbit IgG (3). Immunoprecipitation of S tag protein by 2.5 µg of ab183674. 12% SDS-PAGE. The immunoprecipitated S tag protein was detected by ab183674 diluted at 1/10000. Anti-rabbit IgG was used as a secondary reagent.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors