



Anti-Sall4 antibody [6E3] ab57577

★★★★★ [4 Abreviews](#) [33 References](#) [画像数 6](#)

製品の概要

製品名	Anti-Sall4 antibody [6E3]
製品の詳細	Mouse monoclonal [6E3] to Sall4
由来種	Mouse
アプリケーション	適用あり: WB, IHC-P, Flow Cyt
種交差性	交差種: Human
免疫原	<p>Recombinant fragment corresponding to Human Sall4 aa 954-1054. Sequence:</p> <p>PKEILAPSVNVDPVVWNQYTSMLNGGLAVKTNESVSIQSGGV PTLPVSLG ATSVVNNATVSKMDGSQSGISADVEKPSATDGVPHQFPHFL EENKIAVS</p> <p>Database link: Q9UJQ4</p> <p> Run BLAST with  Run BLAST with</p>
ポジティブ・コントロール	WB: HeLa cell lysate, IHC-P: human testis. Flow Cyt: HeLa cells.
特記事項	<p>This product was changed from ascites to tissue culture supernatant on 12th Feb 2019. Please note that the dilutions may need to be adjusted accordingly. If you have any questions, please do not hesitate to contact our scientific support team.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

バッファー	pH: 7.40 Constituent: 100% PBS
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	6E3
アイソタイプ	IgG1
軽鎖の種類	kappa

アプリケーション


The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab57577の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Predicted molecular weight: 112 kDa.
IHC-P	★★★★★ (4)	Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

ターゲット情報

機能	Probable transcription factor.
組織特異性	Expressed in testis.
関連疾患	Defects in SALL4 are the cause of Duane-radial ray syndrome (DRRS) [MIM:607323]; also known as Okihiro syndrome. DRRS is a disorder characterized by the association of forearm malformations with Duane retraction syndrome. Defects in SALL4 are the cause of oculotoradial syndrome (OORS) [MIM:147750]. Oculotoradial syndrome is an autosomal dominant condition characterized by upper limbs anomalies (radial ray defects, carpal bones fusion), extraocular motor disturbances, congenital bilateral non-progressive mixed hearing loss. Other less consistent malformations include heart involvement, mild thrombocytopenia and leukocytosis (before age 50), shoulder girdle hypoplasia, imperforate anus, kidney malrotation or rectovaginal fistula. The VIC syndrome is an allelic disorder of Duane-radial ray syndrome (DRRS) with a similar phenotype.
配列類似性	Belongs to the sal C2H2-type zinc-finger protein family. Contains 7 C2H2-type zinc fingers.
細胞内局在	Nucleus.

画像




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sall4 antibody [6E3] (ab57577)

Image from Gassei, Kathrin, et. al. PLoS ONE 8.1 (2013): e53976. doi: 10.1371/journal.pone.0053976. Fig 2A,D. Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of testes from juvenile (PND 7 and PND 14) mice labeling Sall4 with ab57577. Testes were dissected and fixed in 4% PFA overnight at 4°C. Fixed tissues were processed for paraffin embedding and 5 µm serial sections were collected. Sections were deparaffinized in xylene (2×15 min), rehydrated in a graded ethanol series (2×100% for 10 min, 1×95% for 5 min, 1×80% for 5 min, 1×70% for 5 min, 1×50% for 5 min, 1×25% for 5 min) and rinsed in 1× DPBS. Slides were then incubated in sodium citrate antigen retrieval buffer (10 mM Sodium Citrate, 0.05% Tween-20, pH 6) for 30 min at 97.5°C and allowed to cool to room temperature. After rinsing twice in 1× DPBS containing 0.1% Tween-20 (DBPS-T), unspecific binding sites in tissue sections were saturated by incubation with blocking buffer (1× DPBS containing 3% bovine serum albumin, 0.1% Triton X-100 and 5% normal serum from the host species of the secondary antibody) for 30 min at room temperature. Primary antibodies were diluted in blocking buffer and added to tissue sections for 90 min at room temperature.

This image was generated using the ascites version of the product.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sall4 antibody [6E3] (ab57577)

This image is courtesy of an Abreview from Zachary Yu-Ching Lin.

IHC-P image of Sall4 staining with ab57577 on tissue sections from adult mouse testis. The sections were subjected to heat-mediated antigen retrieval using Dako antigen retrieval solution. The sections were then blocked with 5% milk for 30 minutes at 25°C, before incubation with ab57577 (1/100 dilution) for 18 hours at 4°C. The secondary was an Alexa-Fluor 488 conjugated goat anti-mouse polyclonal, used at a 1/500 dilution.


This image was generated using the ascites version of the product.



Western blot - Anti-Sall4 antibody [6E3] (ab57577)

Sall4 antibody (ab57577) at 1ug/lane + HeLa cell lysate at 25ug/lane.


This image was generated using the ascites version of the product.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sall4 antibody [6E3] (ab57577)

Sall4 antibody (ab57577) used in immunohistochemistry at 5ug/ml on formalin fixed and paraffin embedded human testis.


This image was generated using the ascites version of the product.



Flow Cytometry - Anti-Sall4 antibody [6E3] (ab57577)

Overlay histogram showing HeLa cells stained with ab57577 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab57577, 1μg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] ([ab91353](#), 2μg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in HeLa cells fixed with 4% paraformaldehyde/permeabilized in 0.1% PBS-Tween used under the same conditions.

This image was generated using the ascites version of the product.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sall4 antibody [6E3] (ab57577)

This image is courtesy of an Abreview from Zachary Yu-Ching Lin.

IHC-P image of Sall4 staining with ab57577 on tissue sections from adult marmoset testis. The sections were subjected to heat-mediated antigen retrieval using Dako antigen retrieval solution. The sections were then blocked with 5% milk for 30 minutes at 25°C, before incubation with ab57577 (1/100 dilution) for 18 hours at 4°C. The secondary was an Alexa-Fluor 488 conjugated goat anti-mouse polyclonal, used at a 1/500 dilution.

This image was generated using the ascites version of the product.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors