

### Anti-SED antibody ab15900

#### 製品の概要

---

製品名	Anti-SED antibody
製品の詳細	Rabbit polyclonal to SED
由来種	Rabbit
特異性	Absorbance readings (410 nm) of less than 0.100 for 10 ng/ml preparations of staphylococcal enterotoxins A through E (excluding D), ET, TSST, and alpha hemolysin.
アプリケーション	<b>適用あり:</b> WB, ELISA
種交差性	<b>交差種:</b> Staphylococcus aureus
免疫原	Full length native protein (purified) corresponding to SED.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

---

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Constituents: 0.0268% PBS, 0.9% Sodium chloride
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

#### アプリケーション

---

**The Abpromise guarantee**      **Abpromise保証は、次のテスト済みアプリケーションにおけるab15900の使用に適用されます**  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 30 kDa.
ELISA		Use a concentration of 10 µg/ml.

## ターゲット情報

**関連性** Staphylococcal enterotoxins represent a group of proteins, which are secreted by *Staphylococcus aureus* and cause the intoxication staphylococcal food poisoning syndrome. The illness is characterised by high fever, hypotension, diarrhea, shock, and in some cases death. Their molecular masses range between 27 and 30 kDa. At present, seven enterotoxins are known, namely A, B, C (subtypes C1, C2, C3), D and E. Their amino acid sequences have been determined and it was shown that all are single chain polypeptides containing one disulfide bond formed by two half cystines located in the middle of the polypeptide chain, which form the so called cysteine loop. Enterotoxins are known to be most potent T cell mitogens. T cell activation accompanied by induction of interleukin 2 and interferon is conditioned by high affinity interaction of S.enterotoxins with class II main histocompatibility complex (MHC) molecules and subsequent presentation of the complex formed to a variable region of the T cell receptor.

**細胞内局在** Secreted

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors