

Anti-Selenium Binding Protein 1/SBP antibody ab90135

★★★★☆ [2 Abreviews](#) [6 References](#) [画像数 6](#)

製品の概要

製品名	Anti-Selenium Binding Protein 1/SBP antibody
製品の詳細	Rabbit polyclonal to Selenium Binding Protein 1/SBP
由来種	Rabbit
アプリケーション	適用あり: WB, ICC/IF, IP, IHC-P
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide corresponding to Human Selenium Binding Protein 1/SBP aa 1-100 conjugated to keyhole limpet haemocyanin. (Peptide available as ab90251)
ポジティブ・コントロール	This antibody gave a positive signal in the following tissue lysates: Human Liver; Human Colon; Human Spleen; Mouse Liver; Mouse Lung; Rat Liver.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS
精製度	Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help. Immunogen affinity purified

ポリモノ
アイソタイプ

ポリクローナル
IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab90135の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (1)	Use a concentration of 1 µg/ml. Detects a band of approximately 52 kDa (predicted molecular weight: 52 kDa).
ICC/IF		Use a concentration of 5 µg/ml.
IP		Use a concentration of 5 µg/ml.
IHC-P	★★★★★ (1)	Use a concentration of 5 µg/ml.

ターゲット情報

機能

Selenium-binding protein which may be involved in the sensing of reactive xenobiotics in the cytoplasm. May be involved in intra-Golgi protein transport.

組織特異性

Highly expressed in liver, lung, colon, prostate, kidney and pancreas. In brain, present both in neurons and glia (at protein level). Down-regulated in lung adenocarcinoma, colorectal carcinoma and ovarian cancer. Two-fold up-regulated in brain and blood from schizophrenia patients.

配列類似性

Belongs to the selenium-binding protein family.


翻訳後修飾

Phosphorylated.
The N-terminus is blocked.

細胞内局在

Nucleus. Cytoplasm > cytosol. Membrane. May associate with Golgi membrane. May associate with the membrane of autophagosomes.

画像



Western blot - Anti-Selenium Binding Protein 1/SBP antibody (ab90135)

All lanes : Anti-Selenium Binding Protein 1/SBP antibody (ab90135) at 1/500 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : SELENBP1 knockout HeLa cell lysate

Lane 3 : HT-29 cell lysate

Lane 4 : SW 480 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

Lanes 1-3 : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) at 1/10000 dilution


Lane 4 : **ab216773** at 1/10000 dilution

Predicted band size: 52 kDa

Observed band size: 60 kDa

Lanes 1-3: Merged signal (red and green). Green - ab90135 observed at 60 kDa. Red - loading control **ab8245** observed at 36 kDa.

ab90135 Anti-Selenium Binding Protein 1/SBP antibody was shown to specifically react with Selenium Binding Protein 1/SBP in wild-type HeLa cells. Loss of signal was observed when knockout cell line **ab265279** (knockout cell lysate **ab257662**) was used. Wild-type and Selenium Binding Protein 1/SBP knockout samples were subjected to SDS-PAGE. ab90135 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 500 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-Selenium Binding Protein 1/SBP antibody (ab90135)

All lanes : Anti-Selenium Binding Protein 1/SBP antibody (ab90135) at 1 µg/ml

Lane 1 : Human liver tissue lysate - total protein ([ab29889](#))

Lane 2 : Human colon tissue lysate - total protein ([ab30051](#))

Lane 3 : Human spleen tissue lysate - total protein ([ab29699](#))

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 52 kDa

Observed band size: 52 kDa


Additional bands at: 50 kDa. We are unsure as to the identity of these extra bands.

Exposure time: 1 minute



Immunocytochemistry/ Immunofluorescence - Anti-Selenium Binding Protein 1/SBP antibody (ab90135)


ICC/IF image of ab90135 stained MCF-7 cells. The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal Goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab90135, 5µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 Goat anti-Rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM. This antibody also gave a positive result in 4% PFA fixed (10 min) HeLa cells at 5µg/ml, and in 100% Methanol fixed (5 min) HeLa, and Hek293 cells at 5µg/ml.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Selenium Binding Protein 1/SBP antibody (ab90135)

IHC image of ab90135 staining in Normal Human Liver formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab90135, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Western blot - Anti-Selenium Binding Protein 1/SBP antibody (ab90135)

All lanes : Anti-Selenium Binding Protein 1/SBP antibody (ab90135) at 1 µg/ml

Lane 1 : Liver (Mouse) Tissue Lysate

Lane 2 : Lung (Mouse) Tissue Lysate

Lane 3 : Liver (Rat) Tissue Lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) preadsorbed (**ab97080**) at 1/5000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 52 kDa

Observed band size: 52 kDa

Exposure time: 1 minute



Selenium Binding Protein 1/SBP was immunoprecipitated using 0.5mg Mouse Liver tissue lysate, 5µg of Rabbit polyclonal to Selenium Binding Protein 1/SBP and 50µl of protein G magnetic beads (+). No antibody was added to the control (-).

The antibody was incubated under agitation with Protein G beads for 10min, Mouse Liver tissue lysate lysate diluted in RIPA buffer was added to each sample and incubated for a further 10min under agitation.

Proteins were eluted by addition of 40µl SDS loading buffer and incubated for 10min at 70°C; 10µl of each sample was separated on a SDS PAGE gel, transferred to a nitrocellulose membrane, blocked with 5% BSA and probed with ab90135.

Secondary: Mouse monoclonal [SB62a] Secondary Antibody to Rabbit IgG light chain (HRP) ([ab99697](#)).

Band: 52kDa; Selenium Binding Protein 1/SBP

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors