

Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] ab62322

リコンビナント **RabMAb**

25 References **画像数 6**

製品の概要

製品名	Anti-SHP2 (phospho Y542) antibody [EP508(2)Y]
製品の詳細	Rabbit monoclonal [EP508(2)Y] to SHP2 (phospho Y542)
由来種	Rabbit
アプリケーション	適用あり: ICC/IF, WB, IP 適用なし: Flow Cyt
種交差性	交差種: Mouse, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	Treated NIH/3T3 cell lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Rat: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EP508(2)Y

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab62322の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		1/100 - 1/250.
WB		1/1000. Detects a band of approximately 68 kDa (predicted molecular weight: 68 kDa). For unpurified use at 1/50000- 1/100000.
IP		1/30. For unpurified use at 1/40

追加情報

Is unsuitable for Flow Cyt.

ターゲット情報

機能

Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus.

組織特異性

Widely expressed, with highest levels in heart, brain, and skeletal muscle.

関連疾患

Defects in PTPN11 are the cause of LEOPARD syndrome type 1 (LEOPARD1) [MIM:151100]. It is an autosomal dominant disorder allelic with Noonan syndrome. The acronym LEOPARD stands for lentigines, electrocardiographic conduction abnormalities, ocular hypertelorism, pulmonic stenosis, abnormalities of genitalia, retardation of growth, and deafness.

Defects in PTPN11 are the cause of Noonan syndrome type 1 (NS1) [MIM:163950]. Noonan syndrome (NS) is a disorder characterized by dysmorphic facial features, short stature, hypertelorism, cardiac anomalies, deafness, motor delay, and a bleeding diathesis. Some patients with Noonan syndrome type 1 develop multiple giant cell lesions of the jaw or other bony or soft tissues, which are classified as pigmented villomoduolar synovitis (PVNS) when occurring in the jaw or joints. Note=Mutations in PTPN11 account for more than 50% of the cases. Rarely, NS is associated with juvenile myelomonocytic leukemia (JMML). NS1 inheritance is autosomal dominant.

Defects in PTPN11 are a cause of juvenile myelomonocytic leukemia (JMML) [MIM:607785]. JMML is a pediatric myelodysplastic syndrome that constitutes approximately 30% of childhood cases of myelodysplastic syndrome (MDS) and 2% of leukemia. It is characterized by leukocytosis with tissue infiltration and in vitro hypersensitivity of myeloid progenitors to granulocyte-macrophage colony stimulating factor.

Defects in PTPN11 are a cause of metachondromatosis (MC) [MIM:156250]. It is a skeletal disorder with radiologic fetarures of both multiple exostoses and Ollier disease, characterized by the presence of multiple enchondromas and osteochondroma-like lesions.

配列類似性

Belongs to the protein-tyrosine phosphatase family. Non-receptor class 2 subfamily.
Contains 2 SH2 domains.
Contains 1 tyrosine-protein phosphatase domain.

ドメイン

The SH2 domains repress phosphatase activity. Binding of these domains to phosphotyrosine-containing proteins relieves this auto-inhibition, possibly by inducing a conformational change in

the enzyme.

翻訳後修飾

Phosphorylated on Tyr-546 and Tyr-584 upon receptor protein tyrosine kinase activation; which creates a binding site for GRB2 and other SH2-containing proteins.

細胞内局在

Cytoplasm.

画像



Immunoprecipitation - Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322)

ab62322 (purified) at 1:30 dilution (2µg) immunoprecipitating SHP2 in NIH/3T3 treated with 50ng/ml PDGF for 40min whole cell lysate.


Lane 1 (input): NIH/3T3 (Mouse embryonic fibroblast) treated with 50ng/ml PDGF for 40min whole cell lysate 10µg

Lane 2 (+): ab62322 & NIH/3T3 treated with 50ng/ml PDGF for 40min whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab62322 in NIH/3T3 treated with 50ng/ml PDGF for 40min whole cell lysate

For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDm/TBST.



Western blot - Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322)

All lanes : Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322) at 1/1000 dilution (Purified)

Lane 1 : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysates

Lane 2 : NIH/3T3 (Mouse embryonic fibroblast) treated with 40ng/ml PDGF for 40 minutes whole cell lysates

Lane 3 : NIH/3T3 (Mouse embryonic fibroblast) treated with 40ng/ml PDGF for 40 minutes then incubated with phosphatase


Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 68 kDa


Observed band size: 68 kDa



Immunocytochemistry/ Immunofluorescence - Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322)

Ab62322 staining SHP2 in THP-1 cells (Human monocytic leukaemia cell line) by ICC/IF (immunocytochemistry/immunofluorescence). Cells were fixed with 4% Paraformaldehyde and permeabilized with 0.1% TritonX-100. Samples were incubated with purified ab62322 at 8.8 µg/ml. Secondary antibody used was AlexaFluor®488 Goat anti-Rabbit (**ab150077**) at 2 µg/ml. Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594)(**ab195889**) used as counterstain at 2.5 µg/ml . DAPI was used as nuclear counterstain. Confocal image showing the expression was increased after treatment with TPA 200nM for 24h and PDGF 50ng/ml for 30min, the signal decreased after treatment with Lambda Protein Phosphatase 31I for 2h.

This image was generated using the unpurified version of the product.



Western blot - Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322)

All lanes : Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322) at 1/200 dilution

Lane 1 : THP-1 (Human monocytic leukemia monocyte) whole cell lysates

Lane 2 : THP-1 (Human monocytic leukemia monocyte) treated with Phorbol-12-myristate-13-acetate and platelet-derived growth factor.

Lane 3 : THP-1 (Human monocytic leukemia monocyte) treated with Phorbol-12-myristate-13-acetate and platelet-derived growth factor. Then the membrane was incubated with phosphatase.

Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution


Predicted band size: 68 kDa

Observed band size: 68 kDa

Blocking and diluting buffer used was 5% NFDM/TBST.

Purified batch of ab62322 was used.

This image was generated using the unpurified version of the product.



All lanes : Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322) at 1/50000 dilution

Lane 1 : NIH/3T3 cell lysates; untreated

Lane 2 : NIH/3T3 cell lysates; treated with PDGF

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : goat anti-rabbit HRP conjugated, at 1/2000 dilution

Predicted band size: 68 kDa

Observed band size: 68 kDa

Beta Tubulin has been included as a loading control.

This image was generated using the unpurified version of the product.

Why choose a recombinant antibody?

Research with confidence
Consistent and reproducible results

Long-term and scalable supply
Recombinant technology

Success from the first experiment
Confirmed specificity

Ethical standards compliant
Animal-free production

Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors